

Team reports reliable, highly efficient method for making stem cells

September 18 2014



A new method resulted in a colony of stem cells, glowing green, derived from one adult immune cell. Credit: Laboratory of Matthias Stadtfeld at NYU Langone Medical center

Scientists at NYU Langone Medical Center have found a way to boost



dramatically the efficiency of the process for turning adult cells into socalled pluripotent stem cells by combining three well-known compounds, including vitamin C. Using the new technique in mice, the researchers increased the number of stem cells obtained from adult skin cells by more than 20-fold compared with the standard method. They say their technique is efficient and reliable, and thus should generally accelerate research aimed at using stem cells to generate virtually any tissue. Stem cells are immature or uncommitted cells that are theoretically capable of becoming any cell type.

"This big boost in efficiency gives us an opportunity now to study stem cell programming mechanisms at high resolution," says Matthias Stadtfeld, PhD, assistant professor of cell biology and a member of the Skirball Institute of Biomolecular Medicine and the Helen L. and Martin S. Kimmel Center for Stem Cell Biology at NYU Langone Medical Center, who led the research.

"This is a very exciting advance," says Ruth Lehmann, PhD, director of the Kimmel Center for Stem Cell Biology and the Skirball Institute at NYU Langone and chair of the Department of Cell Biology. "The new technology developed by the Stadtfeld lab to reprogram <u>differentiated</u> <u>cells</u> efficiently and effectively brings the prospect of stem cell technology for safe use in regenerative medicine ever so much closer."

The standard method for reprogramming skin, blood, or other tissuespecific cell types into "induced <u>pluripotent stem cells</u>" (iPSCs) was reported in 2006 by the laboratory of Kyoto University's Shinya Yamanaka, who later won a Nobel Prize for the achievement. The method involves the artificial expression of four key genes dubbed OKSM (for Oct4, Klf4, Sox2 and myc) whose collective activity slowly prods cells into an immature state much like that of an early embryonic cell.



In principle, one could take a sample of cells from a person, induce the cells to become iPSCs, then multiply the iPSCs in a lab dish and stimulate them to mature towards desired adult cell types such as blood, brain or heart—which then could be used to replace injured or diseased tissue in that same individual.

But there are many formidable technical obstacles, among which is the low efficiency of currently used protocols. Converting most <u>cell types</u> into stable iPSCs occurs at rates of 1 percent or less, and the process can take weeks.

Researchers throughout the world have been searching for ways to boost this efficiency, and in some cases have reported significant gains. These procedures, however, often alter vital cellular genes, which may cause problems for potential therapies. For the new study, reported online today in *Stem Cell Reports*, Dr. Stadtfeld and his laboratory team decided to take a less invasive approach and investigate chemical compounds that transiently modulate enzymes that are present in most cells. "We especially wanted to know if these compounds could be combined to obtain stem cells at high efficiency," Dr. Stadtfeld says.

Two of these compounds influence well known signaling pathways, called Wnt and TGF- β , which regulate multiple growth-related processes in cells. The third is vitamin C (also known as ascorbic acid). Best known as a powerful antioxidant, the vitamin was recently discovered to assist in iPSC induction by activating enzymes that remodel chromatin—the spiral scaffold for DNA—to regulate gene expression.

Simon Vidal, a graduate student in the Stadtfeld lab, and Bhishma Amlani, a postdoctoral researcher, looked first at mouse skin fibroblasts, the most common cell type used for iPSC research. Adding to fibroblasts engineered to express OKSM either vitamin C, a compound to activate Wnt signaling, or a compound to inhibit TGF- β signaling increased iPSC-



induction efficiency weakly to about 1% after a week of cell culture. Combining any two worked a bit better. But combining all three brought the efficiency to about 80 percent in the same period of time.

In another series of experiments the team worked with blood progenitor cells, which usually replace blood cells lost after injury or infection. The OKSM method on its own can slowly convert these cells to stem cells with up to 30 percent efficiency. Using OKSM together with the three compounds brought the efficiency to nearly 100 percent in less than a week. The researchers also achieved nearly 100-percent yield in mouse liver progenitor cells.

Dr. Stadtfeld expects that these dramatic increases in conversion rates of <u>adult cells</u> into embryonic-like stem cells will facilitate future studies of the iPSC induction process, simply by making that induction a more predictable event. "It's just a lot easier this way to study the mechanisms that govern reprogramming, as well as detect any undesired features that might develop in iPSCs," he said.

Vitamin C and the two compounds used to manipulate the Wnt and TGF- β pathways are widely studied and considered to have few unknown or hazardous effects, the researchers said. By contrast, the use of OKSM has in some cases caused undesired features in iPSCs, such as developmental defects. By making iPSC induction more rapid and efficient, though, Dr. Stadtfeld's new technique might also make the resulting stem cells safer. "Conceivably it reduces the risk of abnormalities by smoothening out the reprogramming process," Dr. Stadtfeld says. "That's one of the issues we're following up."

Provided by New York University School of Medicine

Citation: Team reports reliable, highly efficient method for making stem cells (2014, September



18) retrieved 27 June 2024 from <u>https://phys.org/news/2014-09-team-reliable-highly-efficient-method.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.