

## A new approach to on-chip quantum computing

October 2 2014



Cross-polarized pump photons (red and blue) interact in the micro-ring resonator to directly generate cross-polarized correlated photons (green and yellow). Credit: Lucia Caspani

Commercial devices capable of encrypting information in unbreakable codes exist today, thanks to recent quantum optics advances, especially the generation of photon pairs—tiny entangled particles of light. Now, an international team of researchers led by professor Roberto Morandotti of INRS-EMT in Canada, is introducing a new method to



achieve a different type of photon pair source that fits into the tiny space of a computer chip.

The team's method, which generates "mixed up" photon pairs from devices that are less than one square millimeter in area, could form the core of the next-generation of quantum <u>optical communication</u> and computing technology. The research will be presented at The Optical Society's (OSA) 98th Annual Meeting, Frontiers in Optics, being held Oct. 19-23 in Tucson, Arizona, USA.

One of the properties of light exploited within <u>quantum optics</u> is "<u>photon</u> <u>polarization</u>," which is essentially the direction in which the electric field associated with the photon oscillates. The research team set out to find a way to directly "mix up," or cross-polarize, the photons via a nonlinear optical process on a chip.

"While several efforts have been devoted to develop on-chip sources of polarization-entangled photons, the process typically used to generate these photons only allows the generation of photons with the same polarization as the laser beam used to pump the device—either both horizontal or vertical—after which entanglement can be achieved by accurately mixing these states. Now, we have found a way to directly generate cross-polarized photon pairs," says Lucia Caspani, a postdoctoral fellow at INRS-EMT and co-author of the Frontiers in Optics paper.

To generate the cross-polarized photons, Caspani and colleagues used two different laser beams at different wavelengths —one vertically polarized and another horizontally polarized. The approach, however, came with a potential pitfall: the classical process between the two pump beams could destroy the <u>photons</u>' fragile quantum state.

To address this challenge, the team, which also includes researchers



from RMIT University in Australia and City University of Hong Kong, pioneered a new approach based on a micro-ring resonator—a tiny optical cavity with a diameter on the order of tens to hundreds of micrometers—that operates in such a way that energy conservation constraints suppress classical effects while amplifying quantum processes.

While a similar suppression of classical effects has been observed in gas vapors and complex micro-structured fibers, this is the first time it has been reported on a chip, thus opening a clear route for building scalable integrated devices.

"Our approach opens the door to directly mixing different polarizations on a chip," Caspani points out. "At very low power, our device directly generates photon pairs with orthogonal polarizations, which can be exploited for quantum communication and computing protocols."

The fabrication process of the chip is also compatible with that currently used for electronic chips. "It enables a future coexistence of our device with standard integrated circuits," says Caspani, which is a fundamental requirement for the widespread adoption of optical quantum technologies.

**More information:** Presentation FTu2A.2, "Direct Generation of Orthogonally Polarized Photon Pairs via Spontaneous Non-Degenerate FWM on a Chip," takes place Tuesday, Oct. 21 at 11 a.m. MST at the Arizona Ballroom, Salon 8 at the JW Marriott Tucson Starr Pass Resort in Tucson. <u>www.frontiersinoptics.org/</u>

Provided by Optical Society of America



Citation: A new approach to on-chip quantum computing (2014, October 2) retrieved 3 May 2024 from <u>https://phys.org/news/2014-10-approach-on-chip-quantum.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.