

## Lutetia's dark side hosts hidden crater

October 8 2014



Tracing Lutetia's grooves. Credit: ESA/Rosetta/MPS for OSIRIS Team MPS/UPD/LAM/IAA/SSO/INTA/UPM/DASP/IDA

Grooves found on Lutetia, an asteroid encountered by ESA's Rosetta spacecraft, point to the existence of a large impact crater on the unseen side of the rocky world.

Rosetta flew past Lutetia at a distance of 3168 km in July 2010, en route to its 2014 rendezvous with its target comet.

The spacecraft took images of the 100 km-wide asteroid for about two hours during the flyby, revealing numerous impact craters and hundreds



of grooves all over the surface.

Impact craters are commonly seen on all Solar System worlds with solid surfaces, recording an intense history of collisions between bodies. However, grooves are much less prevalent. To date, they have been discovered by visiting spacecraft only on the martian moon Phobos and the asteroids Eros and Vesta.

The way in which grooves are formed on these bodies is still widely debated, but it likely involves impacts. Shock waves from the impact travel through the interior of a small, porous body and fracture the surface to form the grooves.

"For Lutetia, by assuming that the grooves were formed in concentric patterns around their source impact crater, we identified 200 such features falling into distinct 'families', correlated with three different <u>impact craters</u>," describes Sebastien Besse, a research fellow at ESA's Technical Centre, ESTEC, in the Netherlands, and lead author of the paper published inPlanetary and Space Sciencethis month.

One of the groove systems on Lutetia is associated with the Massilia crater and another with the North Pole Crater Cluster, which comprises a number of superimposed craters. Both are on the asteroid's northern hemisphere.





This anaglyph 3D image of Lutetia can be viewed using stereoscopic glasses with red–green or red–blue filters. The two images making up this image were taken several minutes before Rosetta's closest approach to the asteroid on 10 July 2010. The left-eye view was captured at 15:41:39 GMT from a distance of 4274 km from Lutetia's surface and the right-eye view at 15:41:03 GMT from 4038 km (closest approach was at 15:45 GMT). Credit: ESA/H. Sierks (MPS, Göttingen, Germany)

But another group of grooves points to a crater not seen during Rosetta's brief flyby, in the asteroid's <u>southern hemisphere</u>.



Its implied presence has earned it the nickname 'Suspicio'. The grooves related to Suspicio cover a large area on the asteroid, suggesting it may span several tens of kilometres . By comparison, Massilia, the largest known crater on Lutetia, is about 55 km wide, and the largest of the polar cluster is about 34 km across.

"These three major impacts seriously deformed Lutetia's surface," adds Sebastien.



Looking face on at the North Pole Crater Cluster (purple outline) on asteroid Lutetia, with Massilia to the lower left (red outline). Marked on the image are the concentric grooves or 'lineaments' associated with the large craters. The lineaments coloured blue infer the presence of a large crater – nicknamed Suspicio – on the unseen portion of Lutetia. Yellow denotes lineaments not associated with any of the craters discussed in this study. Credit:



ESA/Rosetta/MPS for OSIRIS Team MPS/UPD/LAM/IAA/SSO/INTA/UPM/DASP/IDA

"As with grooves seen on other asteroids that may also be associated with impact events, this study provides new insights into the catastrophic history of these small bodies."

By observing how subsequent small craters lie over the grooves on Lutetia, the scientists determined the relative ages of the three larger cratering events. Massilia is thought be the oldest of the three craters and the polar cluster the youngest, with Suspicio between.

The authors also looked at other, independent measurements of Lutetia, including ground-based observations with the Infrared Telescope Facility and space-based observations with ESA's Herschel and NASA's Spitzer.



The inferred location of Suspicio crater on the unseen southern hemisphere of asteroid Lutetia (marked in blue). The hidden crater could be up to 45 km in



diameter – the blue outlines correspond to diameter estimates of 15, 30 and 45 km, respectively. The crater is inferred based on the numerous grooves or 'lineaments' seen concentric to the crater in the northern hemisphere of the asteroid. There are no image data available for this side of the asteroid, as can be inferred from the blank shape model. Credit: ESA/Rosetta/MPS for OSIRIS Team MPS/UPD/LAM/IAA/SSO/INTA/UPM/DASP/IDA

Shape models derived by Herschel and Spitzer before Rosetta's flyby had already predicted a large depression at the location of Suspicio. The Infrared Telescope Facility suggested different compositions between the northern and southern hemisphere of the asteroid.

Sebastien and his colleagues propose that a large impact, presumably the one forming Suspicio, excavated enough material of a different composition to account for the observed differences.

"Our study ties together several independent analyses of Lutetia into one coherent story that is consistent with the presence of a large impact crater on the far side of the asteroid," says co-author Michael Küppers, from ESA's Space Astronomy Centre in Spain.

"Four years on and we are delighted still to be learning from just two hours' worth of data collected during the Lutetia flyby," says Matt Taylor, ESA's Rosetta project scientist.

"Rosetta is now in its main mission phase at its comet, where we are on the cusp of fantastic results. Rosetta is a true small bodies mission, two asteroids and one comet in single trip."

**More information:** S. Besse, M. Küppers, O.S. Barnouin, N. Thomas, J. Benkhoff, "Lutetia's lineaments," *Planetary and Space Science*,



Volume 101, 15 October 2014, Pages 186-195, ISSN 0032-0633, dx.doi.org/10.1016/j.pss.2014.07.007

## Provided by European Space Agency

Citation: Lutetia's dark side hosts hidden crater (2014, October 8) retrieved 30 April 2024 from https://phys.org/news/2014-10-lutetia-dark-side-hosts-hidden.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.