


Tracking a gigantic sunspot across the Sun

October 31 2014



Super sunspot AR2192 produced 10 significant solar flare while traversing the Earth-side of the sun; six X-class and four above M5-class. Credit: NASA/SDO

An active region on the sun – an area of intense and complex magnetic fields – rotated into view on Oct. 18, 2014. Labeled AR 12192, it soon grew into the largest such region in 24 years, and fired off 10 sizable

solar flares as it traversed across the face of the sun. The region was so large it could be seen without a telescope for those looking at the sun with eclipse glasses, as many did during a partial eclipse of the sun on Oct. 23.

"Despite all the flares, this region did not produce any significant [coronal mass ejections](#)," said Alex Young a solar scientist at NASA's Goddard Space Flight Center in Greenbelt, Maryland. Coronal mass ejections, or CMEs, are giant clouds of solar particles that can affect technology when they reach near-Earth space. "You certainly can have flares without CMEs and vice versa, but most big flares do have CMEs. So we're learning that a big active region doesn't always equal the biggest events."


Such active regions are measured in millionths of a solar hemisphere, where 1 micro-hemisphere, or MH, is about 600,000 square miles. This region topped out at 2,750 MH, making it the 33rd largest region out of approximately 32,000 active regions that have been tracked and measured since 1874. It is the largest sunspot seen since AR 6368, which measured 3,080 MH on Nov. 18, 1990.

The largest five active regions ever observed were between 4,000 and more than 6,000 MH and they all appeared between 1946 and 1951.

On the other hand, the region that produced one of the biggest [solar flares](#) of all time on Sep. 1, 1859 – in what's known as the Carrington event – wasn't even one of the top 50 at only 2,300 MH.

During its trip across the front of the sun, AR 12192 produced six X-class flares, which are the largest flares, and four strong M-class flares. M-class flares are one tenth as strong as X-class flares. The number provides more information about its strength. An M2 is twice as intense as an M1, an M3 is three times as intense, etc.

"Having so many similar flares from the same [active region](#) will be a nice case study for people who work on predicting solar flares," said Dean Pesnell, project scientist for NASA's Solar Dynamics Observatory at Goddard. "This is important for one day improving the nation's ability to forecast space weather and protect technology and astronauts in space."



The largest sunspot since November 1990 is seen traveling across the front of the sun in these images from NASA's SDO, captured Oct. 17-Oct 29, 2014.
Credit: NASA/SDO

The dates and peak times in EDT of the large solar flares from AR 12192 are as follows:

Oct. 19, 1:01 am: X1.1

Oct. 21, 9:59 pm: M8.7

Oct. 22, 10:28 am: X1.6

Oct. 24, 5:41 pm: X3.1

Oct. 25, 1:08 pm: X1.0

Oct. 26, 6:56 am: X2.0

Oct. 26, 8:34 pm: M7.1

Oct. 27, 6:09 am: M6.7

Oct. 27, 10:47 am: X2.0

Oct. 28, 11:32 pm: M6.6

AR 12192 rotated onto the far side of the [sun](#) on Oct. 30, 2014, however as it evolves, we may see a new version of it rotating back into view in two weeks.

Provided by NASA's Goddard Space Flight Center

Citation: Tracking a gigantic sunspot across the Sun (2014, October 31) retrieved 30 April 2024 from <https://phys.org/news/2014-10-tracking-gigantic-sunspot-sun.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--