

Sources of the solar wind

November 5 2014



An ultraviolet image of a network of solar jet taken by the IRIS mission. The region of the Sun shown is about 30,000 kilometers on a side. The red circle highlights a network of about ten jets originating from a line of bright spots. Credit: NASA/IRIS



The solar chromosphere is the region between the Sun's surface and its hot, million-degree corona. Within this complex interface zone, only a few thousand kilometers deep, the density of the gas drops by a factor of about one million and the temperature increases from about five thousand to one million kelvin. Almost all of the mechanical energy that drives solar activity is converted into heat and radiation within this interface. Many physical processes shape the intricate system of solar magnetic fields, energetic particles, and radiation that power the corona. The solar wind itself has long been suspected of originating in this region, perhaps in the cooler "coronal holes," but identifying the precise origin sites required high spatial resolution observations.

In the summer of 2013, NASA launched the Interface Region Imaging Spectrograph (IRIS) mission to study this interface region in ultraviolet light; the CfA and its staff contributed its twenty centimeter telescope as well as to its science team. The latest issue of *Science* features IRIS results, and in it CfA astronomers Hui Tian, Ed DeLuca, Steve Cranmer, Leon Golub, Sean McKillop, Katherine Reeves, Patrick McCauley, PaolaTesta, Mark Weber, and Nicholas Murphy join with their colleagues in announcing the detection of intermittent, small-scale jets from the region and their possible contribution to the stellar wind. The paper is one of five in this special issue of Science which they led or coauthored.

The astronomers found unambiguous evidence for these jets that are roughly about 8,000 thousand kilometers long and 300 kilometers wide, moving between about 80 to 250 kilometers per second, with gas at temperatures of at least one hundred thousand kelvin, and coming in bursts lasting between about 20 and 80 seconds. The upward flowing hot gas may provide the heated mass to the <u>solar wind</u>. The scientists calculate that although intermittent, the ongoing presence of these jets



could provide in principle more than enough material to supply the wind. They caution, however, that much work remains to confirm the conclusions and refine the analyses and models – but in any case the newly discover jet phenomenon will need to be incorporated in any future, detailed wind models.

More information: "Prevalence of small-scale jets from the networks of the solar transition region and chromosphere," H. Tian, E. E. DeLuca, S. R. Cranmer, B. De Pontieu, H. Peter, J. Martínez-Sykora, L. Golub, S. McKillop, K. K. Reeves, M. P. Miralles, P. McCauley, S. Saar, P. Testa, M. Weber, N. Murphy, J. Lemen, A. Title, P. Boerner, N. Hurlburt, T. D. Tarbell, J. P. Wuelser, L. Kleint, C. Kankelborg, S. Jaeggli, M. Carlsson, V. Hansteen, S. W. McIntosh, *Science*, 346, 1255711, 2014.

Provided by Harvard-Smithsonian Center for Astrophysics

Citation: Sources of the solar wind (2014, November 5) retrieved 2 May 2024 from <u>https://phys.org/news/2014-11-sources-solar.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.