

Barren deserts can host complex ecosystems in their soils

December 22 2014, by Adam Hadhazy, Astrobio.net



A biological soil crust in Hovenweep National Monument, a region located in southwestern Colorado and southeastern Utah. Credit: Nationalparks (Transferred by Nihonjoe)/Wikipedia

"Biological soil crusts" don't look like much. In fact, people often trample right over these dark, or green-tinted, sometimes raised patches in the desert soil. But these scruffy stretches can house delicate

ecosystems as varied and complexly interwoven as that of a lush, tropical rainforest.

Life forms including bacteria, algae, fungi and lichens, as well as plants such as mosses and liverworts, can band together to create biological soil crusts in dry, nutrient-starved environments. Scientists are just beginning to document the diversity of species that call biological soil crusts home.

"These are incredibly diverse microbial communities with hundreds of different organisms," said Jason Raymond, assistant professor in the School of Earth and Space Exploration at Arizona State University. "If you were counting animals in the Amazon, you wouldn't come close to the diversity of these biological soil crusts."

Raymond is the senior author of three new papers in the scientific journal *Genome Association*, which shed light on the microbes that commonly set up shop in biological soil crusts in Utah's Moab Desert. The papers present a genome of three different bacteria. These genomes contain genes known to enable certain biological forms and functions. Identifying these genes therefore speaks to the interplay of these bacteria as they eke out a living in their shared, severe environment.

Figuring out how life thrives in biological soil crusts, in conditions that would fells most other life on the planet, will help in gauging the habitability of other worlds.

"Biological soil crusts are an outstanding example of nature coping with a really challenging set of environmental conditions," said Raymond. "We're pushing as close as we can to extreme environments on other planets."

Starved in the desert

Although the Moab Desert is hot and dry, the biggest challenge for life in its biological soil crusts, as in other such places, is obtaining nutrients—food, essentially. The new studies reveal the array of adaptive tools the microbes possess for ensnaring scarce, vital nutrients and preventing them from leaching away into the environment.

On a basic level, humans, like all other life, do not need hot dogs to eat or ice tea to drink to survive. When it comes to food and beverage, it boils down to chemical elements. The most basic set of chemical elements life typically needs is summed up by the acronym CHNOPS, made up of the letter signifiers of the elements carbon, hydrogen, nitrogen, oxygen, phosphorus and sulfur. (Complex life, like humans, need a whole bunch of other elements as well, such as selenium and iodine.)

In the Moab Desert, carbon and nitrogen are two particularly dear elements. In more hospitable habitats, like a grassland in a temperate climate, carbon—organic matter—is readily available from fallen leaves, dead plants and animals, and so on. Nitrogen is "fixed," or nabbed from the atmosphere by numerous microbes and taken up by plants for subsequent dissemination in the environment. In the Moab, in contrast, not enough plants or animals can survive to keep these two elements widely available for use by other organisms.

"If you've been there or seen pictures, you know how barren that landscape is," said Raymond. "In terms of fundamental [nutrient availability](#) of carbon and nitrogen, these things are tough to get your hands on when you're in a biological soil crust."

Living on the edge

To get a bead on understanding how the Moab Desert's microbial critters make do, Raymond and his Arizona State University colleagues obtained

samples of biological soil crusts found there. Three microbes grew readily in a growth medium in the lab. The medium, however, was not exactly a smorgasbord. To get by on the available nutrients, the microbes still had to be resourceful and obtain nutrients from the air, such as carbon via carbon dioxide gas, or use one of a small handful of compounds the team supplied in the media.

"It was a very minimal media," said Raymond. "If something's growing on there, you know it's able to fix carbon and nitrogen out of the atmosphere, or using one of the specifically chosen carbon and nitrogen compounds we add to the media. Either way, these organisms are very efficient recyclers of organic material."



A picture of Canyonlands National Park near the city of Moab, Utah. Credit: NOAA/NGDC, John Lockridge, Longmont, Colorado

After growing the cultures, the scientists sequenced the genes of the organisms present. Three different species in the *Microvirga*, *Bacillus* and *Massilia* genera stood out. The three species are alike in some ways. For instance, all have genes for making structures such as flagella—whiplike tails—to allow them to get closer to areas with favorable nutrient availability. Once there, all three microbes have genes for producing "biofilms," a tactic of binding together to remain in place.

Significant differences in their genetic toolkits exist as well. Each bacterium tries to occupy a niche in the biological soil crust community, in partnership or perhaps even at the occasional expense of its neighbors. The *Bacillus* strain, for instance, has genes for pumping out what are known as siderophores. These molecules bind readily to iron, for instance, another chemical element that many organisms need to survive.

The *Microvirga* bacterial strain identified by Raymond and colleagues has genes for sucking up siderophores from the environment, but not for making them. It seems that the *Microvirga* benefits from the *Bacillus* going to the trouble of sending out siderophores.

"The interesting thing is that the *Bacillus* sends out these little shuttles to hopefully get ahold of iron and bring it back, but there is no guarantee," said Raymond.

The *Microvirga*, in this instance, might well be freeloaders.

Letting nothing go to waste


In other ways, the three species coexist peacefully, particularly when there's no speakable limit to the resource. All three benefit from the general ability to obtain nutrients from the air, which is certainly a "more the merrier" type of situation in a biological soil crust.

"There is an incredible efficiency in recycling organic matter so that it doesn't go back into the environment," noted Raymond.

On an individual basis, the bacteria can complement each other. The *Massilia*, for example, fills a niche by apparently being able to survive in oxygen-free, "anaerobic" conditions. That ability suggests it contributes to the overall community's wellbeing by still retaining nutrients in the shared environment in internal or underground locations sealed off from the outside air.

"Each microbe has a specialty and they are complementary," said Raymond.

Characterizing all three [microbes](#), Raymond said: "They're sort of the trash compactors of this community. They'll literally grab ahold of any nutrient-rich sources in the environment and try to metabolize it. That's a testament to how limited this environment is."



A *Bacillus* species with evident flagella, the whiplike tails many microbes use to move around. Credit: CDC/Dr. William A. Clark/Wikipedia

The new studies suggest the high degree to which species living in biological soil crusts rely on their neighbors to play respective parts in nutrient fixation, processing, dead member decomposition, and more.

"Nearly every organism in this community is dependent on some other organism's waste product or byproduct," said Raymond.

Delicate, yet steadfast

Interdependence at such a high level does leave biological soil crusts

vulnerable. Human activity in particular, such as four-wheeling in remote desert environs caked by biological soil crusts, can devastate whole mini-ecosystems, which might take centuries to recover.

"Any perturbation has the potential to have catastrophic consequences," said Raymond.

On the other hand, the ties that bind in biological soil crusts do point, more positively, to how organisms can cooperate to turn a wasteland into an oasis. Environments that scientists would expect to be sterile often astonish us with the lengths to which their denizens go to survive. And as in the case of biological soil crusts, surprising complexity can arise in some of the harshest places.



A moss species, called hairless twisted moss, growing as part of a biological soil crust in Utah. The member species of biological soil crusts each play important roles in maintaining the health of the overall community. Credit: NPS/Neal Herbert

The same paradigm probably holds true for other worlds with conditions

that look abjectly dismal for life.

"One of the things that constantly surprises me is, wherever people seem to go on the planet, there's life," said Raymond. "Anywhere we go on Earth, life has figured out a way to penetrate that niche and take advantage of some aspect of that [environment](#)."

Accordingly, Raymond thinks that because defining habitability is proving so slippery, astrobiologists might want to take the opposite approach in setting out the parameters of clear uninhabitability.



An artist's impression of sunset on an austere exoplanetary landscape, in this case of the super-Earth Gliese 667 Cc. The exoplanet resides in a triple star system and the three suns are visible in the sky. Credit: ESO/L. Calçada

"We're trying to find examples of the most extreme environments we

can on Earth," said Raymond. "We're trying to get into places where biology runs headlong into the inorganic part of the Earth, where any nutrient you want to pick up you've got to get from a rock or the atmosphere."

Essentially, if life can make it there, in these places on Earth, it might just make it anywhere.

This story is republished courtesy of NASA's Astrobiology Magazine. Explore the Earth and beyond at www.astrobio.net.

Source: Astrobio.net

Citation: Barren deserts can host complex ecosystems in their soils (2014, December 22) retrieved 19 April 2024 from <https://phys.org/news/2014-12-barren-host-complex-ecosystems-soils.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--