

Pragmatic approach to saving what can be saved

December 22 2014



Dr. Kaiser-Bunbury (center) and colleagues on the island of Mahé, Seychelles.
Credit: Christopher Kaiser-Bunbury

How can biodiversity be preserved in a world in which traditional ecosystems are increasingly being displaced by "man-made nature"? Biologists at the TU Darmstadt and ETH Zurich have developed a new concept for conservation measures that incorporates current landscapes formerly considered ecologically "of little value."

Numerous experiences from islands have shown that this concept has a positive effect on biodiversity. Now the authors are proposing upscaling these experiences to other landscape scenarios.

In a human-dominated world that contains only little "historical" nature, the term ecosystem can no longer be a synonym for unspoilt nature. The term "novel ecosystems" was coined a few years ago to describe disturbed ecosystems in which biodiversity has been significantly altered as the result of human intervention.

"In our new [conservation](#) framework we argue that this strict distinction between historic and novel ecosystems should be reconsidered to aid conservation," pollination biologist Dr. Christopher Kaiser-Bunbury describes the approach, which is not without controversy.

On continents with vast natural parks, such as the USA and Africa, critics fear that the new concept could weaken the protection of historic nature by, for instance, redirecting financial resources towards more active intervention and design of ecosystems.



A day gecko *Phelsuma ornata* with a blossom *Gastonia mauritiana* on the island of Mauritius. Credit: Christopher Kaiser-Bunbury

The team of Darmstadt and Zurich biologists, however, propagates a reconciling approach. "Our framework combines strategies that were, until now, considered incompatible. Not only historic wildlands are worth protecting, but also designed cultural landscapes. Given the increased anthropogenic pressure on nature, we propose a multi-faceted approach to preserve biodiversity: to protect historic nature where ecologically viable; to actively create new, intensively managed ecosystems; to accept novel ecosystems as natural, wild landscapes; and to convert agricultural and other cultivated landscapes while generally maintaining land-use priorities."

Agricultural landscapes "of little value" belong on the agenda

New ecosystems may also include maize fields and banana plantations, as agricultural land can be used to preserve biodiversity. In fact, necessary measures are relatively easy to implement and comparatively inexpensive. Trials in Europe involving hedges and meadow strips along fields, for example, have shown that many animal species use these areas for feeding and nesting.

Such modifications also create corridors between habitats that are traditionally worth protecting. "The individual measures proposed here are not novel but what is needed is an overall concept that combines these measures on a landscape level. And this is something that has been tested on many oceanic islands – with considerable success."

Lessons from islands



Experiences from islands have shown that the concept has a positive effect on biodiversity. Credit: Christopher Kaiser-Bunbury

The studies by the Darmstadt and Swiss biologists have shown that [biodiversity conservation](#) on regionally heterogeneous islands, such as Galapagos, Hawaii, Fiji or Seychelles, illustrates the successful implementation of such an integrated concept. On the Seychelles, for instance, the combined [conservation measures](#) include the strict protection of natural cloud forest on a few mountain tops, the management of abandoned cinnamon plantations, and green urban areas such as gardens.

The recovery of threatened species and a halt to the decline of native biodiversity are indicators of the success of these conservation strategies. "At the same time, though, we need to know more about how invasive species influence [biodiversity](#)," adds Professor Nico Blüthgen in the light of current investigations by the TU Darmstadt. "For example, native plants on Hawaii have not developed a protective mechanism against immigrant ants and are therefore threatened by their invasion." His working group addresses the consequences of species extinction on ecosystem functioning.

In another study the ecologists aim to investigate how the concept developed on islands can be intensively tested in different landscape settings, including on a larger scale on continents.

More information: "Reconciling conflicting perspectives for biodiversity conservation in the Anthropocene." *Frontiers in Ecology and the Environment* 12: 131–137. [dx.doi.org/10.1890/120201](https://doi.org/10.1890/120201)

Provided by Technische Universität Darmstadt

Citation: Pragmatic approach to saving what can be saved (2014, December 22) retrieved 3 May 2024 from <https://phys.org/news/2014-12-pragmatic-approach.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.