


# A read head for quantum computers? Graphene layer reads optical information from nanodiamonds

December 1 2014

---



Vision of a future quantum computer with chips made of diamond and graphene.  
Credit: Christoph Hohmann, NIM

Nitrogen-vacancy centers in diamonds could be used to construct vital components for quantum computers. But hitherto it has been impossible

to read optically written information from such systems electronically. Using a graphene layer, a team of scientists headed by Professor Alexander Holleitner of the Technische Universität München has now implemented just such a read unit.


Ideally, diamonds consist of pure carbon. But natural diamonds always contain defects. The most researched defects are nitrogen-vacancy centers comprising a [nitrogen atom](#) and a vacancy. These might serve as highly sensitive sensors or as register components for quantum computers. However, until now it has not been possible to extract the optically stored information electronically.

A team headed by Professor Alexander Holleitner, physicist at the TU München and Frank Koppens, physics professor at the Institut de Ciències Fòniques near Barcelona, have now devised just such a methodology for reading the stored information. The technique builds on a direct transfer of energy from nitrogen-vacancy centers in nanodiamonds to a directly neighboring graphene layer.

## **Non-radiative energy transfer**

When laser light shines on a nanodiamond, a light photon raises an electron from its ground state to an excited state in the nitrogen-vacancy center. "The system of the excited electron and the vacated [ground state](#) can be viewed as a dipole," says Professor Alexander Holleitner. "This dipole, in turn, induces another dipole comprising an electron and a vacancy in the neighboring graphene layer."

In contrast to the approximately 100 nanometer large diamonds, in which individual nitrogen-vacancy centers are insulated from each other, the [graphene](#) layer is electrically conducting. Two gold electrodes detect the induced charge, making it electronically measurable.



Laboratory set-up measuring the interaction between graphene and nano-diamonds with implanted nitrogen-vacancy centers. Credit: Astrid Eckert / TUM

### **Picosecond electronic detection**

Essential for this experimental setup is that the measurement is made extremely quickly, because the generated electron-vacancy pairs disappear after only a few billionths of a second. However, the technology developed in Holleitners laboratory allows measurements in the picosecond domain (trillionths of a second). The scientists can thus observe these processes very closely.

"In principle our technology should also work with dye molecules," says doctoral candidate Andreas Brenneis, who carried out the measurements in collaboration with Louis Gaudreau. "A diamond has some 500 point defects, but the methodology is so sensitive that we should be able to

even measure individual dye molecules."

As a result of the extremely fast switching speeds of the nanocircuits developed by the researchers, sensors built using this technology could be used not only to measure extremely fast processes. Integrated into future quantum computers they would allow clock speeds ranging into the terahertz domain.

**More information:** Ultrafast electronic readout of diamond nitrogen-vacancy centres coupled to graphene, Andreas Brenneis, Louis Gaudreau, Max Seifert, Helmut Karl, Martin S. Brandt, Hans Huebl, Jose A. Garrido, Frank H. L. Koppens and Alexander W. Holleitner, *Nature Nanotechnology*, Advanced online publication, December 1, 2014 – [DOI: 10.1038/nnano.2014.276](https://doi.org/10.1038/nnano.2014.276)

Provided by Technical University Munich

Citation: A read head for quantum computers? Graphene layer reads optical information from nanodiamonds (2014, December 1) retrieved 3 May 2024 from <https://phys.org/news/2014-12-quantum-graphene-layer-optical-nanodiamonds.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.