

Peat fire emissions may shed light on climate change

January 16 2015



Rajan Chakrabarty, Ph.D., Washington University in St. Louis. Credit: Washington University in St. Louis

Wildfires, which send hot flames and smoke high into the air, create black carbon emissions associated with climate change and risk to human health. Carbon emissions from wildfires in the contiguous U.S. are expected to increase by 50 percent by 2050 and to double by 2100.

But there's another major yet often overlooked contributor to [carbon emissions](#). Burning of peatlands—an organic mixture of decayed and compacted leaves—are the largest fires on earth. These high-moisture-containing fuels make up nearly three-fourths of earth's land mass and are believed to be the largest emitter of carbon from [wildfires](#) to the atmosphere. However, unlike flaming fires, peatlands smolder underground for weeks and months and emit a whitish smoke. While plenty is known about the environmental effects of black carbon, the effects of peatfire smoke on [human health](#) and the climate are largely unstudied and an emerging field.

To study the climatic effects of carbon-containing aerosols emitted from peat fires, Rajan Chakrabarty, PhD, assistant professor of environmental engineering at Washington University in St. Louis' School of Engineering & Applied Science, has received a three-year, \$530,000 grant from the National Aeronautics and Space Administration (NASA). Brent Williams, PhD, assistant professor of environmental engineering, is a co-investigator on the project, as is Wei Min Hao, PhD, an atmospheric chemist at the USDA Forest Service Fire Sciences Laboratory in Montana.

"This project is going to provide the much-needed information on peat smoke aerosol properties for integration in satellite retrieval algorithms and climate models," Chakrabarty says. "Based on my initial findings, I hypothesize the peat smoke is made up of brown carbon and not black carbon. Brown carbon is a class of organic carbon aerosol which, unlike black carbon, strongly absorbs incoming solar radiation in the shorter wavelengths, or near ultraviolet."

Peat fires are difficult to extinguish, burn at a lower temperature, produce more smoke, generate particles that can be irritating to the eyes and respiratory system, and are among the least understood of fires, according to the U.S. Environmental Protection Agency. In 1997, a

period of extreme haze caused by the spread of smoldering peat fires in Indonesia led to a rapid increase in respiratory illnesses and disrupted shipping and aviation routes, creating an estimated 0.81 to 2.57 billion tons of carbon gases in that region.

With the new grant, Chakrabarty and his collaborators will study the origin of brown carbon particles, their chemical makeup, how many particulates are being emitted and determine their optical properties, or how they behave in light, using a photoacoustic technique, which uses non-invasive imaging based on sound and light.

When scientists and engineers collect the smoke particles on a filter, the residue is yellow to brown in color, giving clues about its properties, Chakrabarty says.

"When you see the yellowish or brown color, it implies that the brown carbon is reflecting all light but the blue violet wavelengths, and that's what it's absorbing," Chakrabarty says. "That has significant implications because we know that black carbon is much less in terms of mass emissions compared to organic carbon. And if organic carbon has absorption right near visible or near ultraviolet light when the solar spectrum starts to peak, it will add further warming on top of [black carbon](#), and we don't know how much."

Chakrabarty and Williams will study laboratory-scale fires from peat fuel types found worldwide at the USDA Fire Science Laboratory combustion chamber using their real-time optical and chemical instruments, which they will transport to Montana in the summer of 2016. They will compile the results into a reference for other scientists, called a look-up table, which will provide researchers and climate modelers with properties for the different fuel types. Since NASA has the largest inventory of fire images from satellite, its scientists will integrate the look-up table into several of its satellite algorithms and

climate models, Chakrabarty said.

Chakrabarty first studied brown carbon particles as a graduate student at the University of Nevada, Reno, in a pilot study at the USDA Fire Science Laboratory. He observed large-scale [brown-carbon](#) aerosol being freshly emitted as primary particles from smoldering combustion of two kinds of duff, or the surface organic layer containing litter, lichen, live and dead moss and organic soil. The combustion of duff is the largest contributor to smoldering smoke production in forest regions in the Northern Hemisphere and contributes between 46 percent and 72 percent of all wildland fire [carbon](#) emissions annually.

Provided by Washington University in St. Louis

Citation: Peat fire emissions may shed light on climate change (2015, January 16) retrieved 20 April 2024 from <https://phys.org/news/2015-01-peat-emissions-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.