

## Do sexually transmitted diseases drive variation in mammalian immunity?

January 23 2015



A young brown rat (Rattus norvegicus) Testes and white blood cells shown. Credit: IZW/Jundong Tian

It has been assumed that the increased transmission of sexuallytransmitted diseases in the case of mating promiscuity is influential in shaping the immune system of mammals. Results published in the scientific journal *Functional Ecology* this week demonstrate that this



simple idea does not apply to rodents, and that living circumstances and the environment can be a key factor in determining variation in immune investment among mammals.

The <u>immune system</u> protects organisms against diseases. Therefore, detecting the factors which shape this system is of great interest to both human and animal medical science. One major question is whether exposure to sexually-transmitted diseases is a main force driving variation in mammalian immunity. Some evidence suggested that this was the case for primates and carnivores and until now was thought to apply to all mammals. Researchers from the German Leibniz Institute for Zoo and Wildlife Research (IZW) have now shown that for rodents, representing 40 % of all living mammal species, this is not the case.

In order to assess the 'sexually-transmitted diseases' hypothesis, scientists from the IZW retrieved data from 145 published studies on rodents. They used the number of total white blood cells and their two main components, neutrophils and lymphocytes, as measures of immunity. They quantified the risk of sexually-transmitted diseases by using measures of testes mass, a known predictor of the number of mating partners across species. The effects of body mass differences and the living circumstances (captive vs. free-ranging) were also taken into consideration in the analyses. The findings demonstrated that large species displayed an increase in the number of immune cells. Also, individuals from captive populations exhibited higher lymphocyte counts than individuals from free-ranging ones. However, testes mass did not vary with immunity, which suggests that sexually-transmitted diseases do not play a major role in shaping the <u>rodent</u> immune system.

"As we know that the prevalence of sexually transmitted pathogens as well as immunological parameters can differ between mammalian groups, pressure by sexually-transmitted diseases may not be the primary determinant of driving immune function in all mammals", says Jundong



Tian, lead author of the study. "Moreover, there is also substantial evidence for effects of captivity upon the physiology of organisms. This suggests that findings derived from captive populations need to be considered very cautiously if we want to understand how evolutionary forces have acted on the immune system over millions of years", comments Gábor Czirják, principal investigator of the study.

Identifying factors which shape immunity in <u>mammals</u> is likely to ultimately lead to advances in medical treatment. Studies applied to rodents are particularly important in this respect, as mice and rats – key representatives of this mammalian group – continue to serve as a major model animal source for biomedical research.

**More information:** Tian JD, Courtiol A, Schneeberger K, Greenwood AD, Czirják GÁ (2014): "Circulating white blood cell counts in captive and wild rodents are influenced by body mass rather than testes mass, a correlate of mating promiscuity." *Functional Ecology*. <u>DOI:</u> <u>10.1111/1365-2435.12394</u>

## Provided by Forschungsverbund Berlin e.V. (FVB)

Citation: Do sexually transmitted diseases drive variation in mammalian immunity? (2015, January 23) retrieved 2 May 2024 from <u>https://phys.org/news/2015-01-sexually-transmitted-diseases-variation-mammalian.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.