

Where did all the stars go? Dark cloud obscures hundreds of background stars

January 7 2015



The Wide Field Imager (WFI) on the MPG/ESO 2.2-metre telescope at the La Silla Observatory in Chile snapped this image of the dark nebula LDN 483. The object is a region of space clogged with gas and dust. These materials are dense enough to effectively eclipse the light of background stars. LDN 483 is located about 700 light-years away in the constellation of Serpens (The Serpent). Credit:

ESO

Some of the stars appear to be missing in this intriguing new ESO image. But the black gap in this glitteringly beautiful starfield is not really a gap, but rather a region of space clogged with gas and dust. This dark cloud is called LDN 483—for Lynds Dark Nebula 483. Such clouds are the birthplaces of future stars. The Wide Field Imager, an instrument mounted on the MPG/ESO 2.2-metre telescope at ESO's La Silla Observatory in Chile, captured this image of LDN 483 and its surroundings.

LDN 483 is located about 700 light-years away in the constellation of Serpens (The Serpent). The cloud contains enough dusty material to completely block the visible light from background [stars](#). Particularly dense molecular clouds, like LDN 483, qualify as dark nebulae because of this obscuring property. The starless nature of LDN 483 and its ilk would suggest that they are sites where stars cannot take root and grow. But in fact the opposite is true: [dark nebulae](#) offer the most fertile environments for eventual star formation.

Astronomers studying [star formation](#) in LDN 483 have discovered some of the youngest observable kinds of baby stars buried in LDN 483's shrouded interior. These gestating stars can be thought of as still being in the womb, having not yet been born as complete, albeit immature, stars.

In this first stage of stellar development, the star-to-be is just a ball of gas and dust contracting under the force of gravity within the surrounding molecular cloud. The protostar is still quite cool—about -250 degrees Celsius—and shines only in long-wavelength submillimetre light. Yet temperature and pressure are beginning to increase in the fledgling star's core.

This earliest period of star growth lasts a mere thousands of years, an astonishingly short amount of time in astronomical terms, given that stars typically live for millions or billions of years. In the following stages, over the course of several million years, the protostar will grow warmer and denser. Its emission will increase in energy along the way, graduating from mainly cold, far-infrared light to near-infrared and finally to [visible light](#). The once-dim protostar will have then become a fully luminous star.

As more and more stars emerge from the inky depths of LDN 483, the dark nebula will disperse further and lose its opacity. The missing background stars that are currently hidden will then come into view—but only after the passage of millions of years, and they will be outshone by the bright young-born stars in the cloud.

Provided by ESO

Citation: Where did all the stars go? Dark cloud obscures hundreds of background stars (2015, January 7) retrieved 28 April 2024 from <https://phys.org/news/2015-01-stars-dark-cloud-obscures-hundreds.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--