


# Autonomous atom assembly of nanostructures using a scanning tunneling microscope

February 16 2015, by Joseph Stroscio

---



Automated assembly of individual cobalt atoms on an atomically flat copper surface into simple geometric shapes, a square, a triangle, and a circle. From left to right, each figure shows the configuration after each atom move. Image size  $15 \text{ nm} \times 15 \text{ nm}$ . Center: Perfect assembly of the NIST logo after four steps of automated assembly. Image size  $40 \text{ nm} \times 17 \text{ nm}$ . All images are shown in colored 3D top view with light shadowing with a height range of  $\approx 100 \text{ pm}$ .

NIST researchers have demonstrated the autonomous computer-controlled assembly of atoms into perfect nanostructures using a low temperature scanning tunneling microscope. The results, published in an invited article in the *Review of Scientific Instruments*, show the construction without human intervention of quantum confined two-dimensional nanostructures using single atoms or single molecules on a copper surface.

A major goal of nanotechnology is to develop so-called "bottom up" technologies to arrange matter at will by placing atoms exactly where one wants them in order to build [nanostructures](#) with specific properties or function. The researchers, led by Robert Celotta and Joseph Stroscio from the CNST, have demonstrated the first steps towards achieving that capability using the atom manipulation mode of a [scanning tunneling microscope](#) (STM) in combination with autonomous motion algorithms.

The team, which includes Stephen Balakirsky (previously in EL and now at Georgia Tech), Aaron Fein (PML), Frank Hess (previously in the CNST), and Gregory Rutter (previously in the CNST and now at Intel), used autonomous algorithms to manipulate single [atoms and molecules](#), much like the algorithms for "hands-free" car driving. The system works by first scanning the locations of available atoms on the surface. It then specifies the desired coordinates of atoms of a nanostructure, and autonomously calculates and directs the trajectories for the STM probe tip to move all the atoms to their desired locations.

The team was able to demonstrate that it could autonomously construct [cobalt atoms](#) into nanostructures that confine the quantum properties of the copper's surface electrons. It then used the STM to measure those properties. In addition to demonstrating the construction of nanostructures made out of atoms, they demonstrated that it was possible to construct nanoscale lattices made of carbon monoxide molecules and to tailor-make interacting quantum dots formed from vacancies in the

carbon monoxide lattices.

The researchers believe that an approach based on autonomous construction of [atoms](#) and molecules using this technique could be the foundation for an easily accessed toolkit for producing tailored quantum states with applications in [quantum information processing](#) and nanophotonics.

**More information:** Autonomous assembly of atomically perfect nanostructures using a scanning tunneling microscope, R. J. Celotta, S. B. Balakirsky, A. P. Fein, F. M. Hess, G. M. Rutter, and J. A. Stroscio, *Review of Scientific Instruments* 85, 121301 (2014).

via [www.nist.gov/cnst/automated\\_atom\\_assembly.cfm](http://www.nist.gov/cnst/automated_atom_assembly.cfm)

Provided by National Institute of Standards and Technology

Citation: Autonomous atom assembly of nanostructures using a scanning tunneling microscope (2015, February 16) retrieved 19 April 2024 from <https://phys.org/news/2015-02-autonomous-atom-nanostructures-scanning-tunneling.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.