

# Killing fish egg fungus with a disinfectant

February 19 2015, by Sandra Avant

---



To control fungus, ARS toxicologist Dave Straus treats channel catfish eggs with peracetic acid. Credit: Cindy Ledbetter

A product used as a disinfectant in agriculture, food preparation, and medical facilities also kills a fungus that causes the disease saprolegniasis on catfish eggs, and it has the potential to treat harmful parasites.

At the Agricultural Research Service's Harry K. Dupree Stuttgart [Arkansas] National Aquaculture Research Center, toxicologist David Straus and his colleagues took a close look at the use of peracetic [acid](#)

—a stabilized mixture of acetic acid (concentrated vinegar) and hydrogen peroxide—for treating catfish eggs. They first determined how much peracetic acid was toxic to hatching catfish fry. The toxicity study indicated that a safe treatment rate was 2.2 parts per million (ppm) peracetic acid for 2-day-old yolk-sac fry and 1.3 ppm for 7-day-old swim-up fry.

"The advantage of using peracetic acid is that it does not produce any residues that would harm fry or the environment," Straus says. "It is safe and effective to use on fish at a low dose, breaks down rapidly into harmless residues, and does not leave an environmental footprint like other chemicals."

Peracetic acid is approved by the U.S. Environmental Protection Agency for many uses in the United States, including disinfecting wastewater and sterilizing items for the food industry and hospitals. It is not approved for aquaculture use in the United States, but it is in Europe. "In Europe, the compound is considered a safe and effective replacement for banned chemicals and antibiotics," Straus adds.

"My colleagues in Germany and Denmark are using it to control fungus on adult fish, and we have also found it very effective for certain parasites," Straus says. "We are working together on acute toxicity studies in different species of fish, because this information is lacking. You need to know how safe it is for fish before you can treat them."



Fungus on untreated channel catfish eggs. Credit: Dave Straus

In a separate study, Straus looked at the effectiveness of different concentrations of peracetic acid in killing fungus on eggs. His experiments included five peracetic acid concentrations (2.5, 5, 10, 15, and 20 ppm) and an untreated control in a flow-through system. Eggs were treated twice daily until the embryos developed eyes. Hatching was complete in 9 days and the fry were counted to determine the survival rate in each treatment.

Fungal growth was severe in the controls, resulting in 11 percent survival compared to 60 percent in the 2.5-ppm group. Very little fungus was present in treatments of 2.5 ppm or higher. However, treatments of 15 and 20 ppm were toxic to eggs. "Although the highest survival rate of hatched fry was at 5 ppm, we found 2.5 ppm to be a safe treatment," Straus says. "This rate is very low, and peracetic acid is flushed out of

the troughs quickly."

Fungal infections such as saprolegniasis in hatchery-reared catfish eggs can cause serious losses of young fish, Straus says. This research provides significant information that could help aquaculture producers deal with fungus problems on eggs.



Healthy, fungus-free channel catfish eggs, less than 24 hours old. Credit: Dave Straus

Provided by Agricultural Research Service

Citation: Killing fish egg fungus with a disinfectant (2015, February 19) retrieved 23 May 2024 from <https://phys.org/news/2015-02-fish-egg-fungus-disinfectant.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.