


Quadruplets in a stellar womb

February 23 2015



An image at radio wavelengths of a young stellar quadruplet. Astronomers have discovered four distinct gas condensations in a clumpy, filamentary gas cloud (white) surrounded by dust (blue). The locations of the condensations in this image are marked with black and red dots. The four condensations are destined to form a bound multiple star system, and one of them (the red dot) has already turned on as a protostar. Credit: Nature; Pineda

More than half of all stars are in multiple systems: binary stars, or even triplets or quadruplets, that orbit one another. No one is quite sure how or why they form, but the effects can be significant, for example

influencing the character of their planets. Our Sun is uncommon in having no companion star, perhaps suggesting that its configuration of planets is equally uncommon.

There are two principal ideas about how multiple stars form: fragmentation in the early stages of birth, or the gravitational capture of a nearby star later on. Computer simulations of [star formation](#) find that both are reasonable possibilities, and so astronomers have been trying to make observations to refine the models and the conclusions. Writing in this week's journal *Nature*, Alyssa Goodman and her collaborators report finding a nearby stellar nursery where quadruplets are being born. The region is in the star forming molecular cloud in the direction of the constellation of Perseus, about 825 light-years away. Scientists have known for decades about a protostar in this area, a [dense core](#) of material that is developing into a small star about one-tenth of a solar-mass in size.

Using radio wavelength observations of dense molecular gas, ammonia in particular, the team discovered that around this protostar are several filamentary gas structures in which they detected three other condensations. The other three embryos are two to three times more massive than the main protostar, and models suggest they will become stars soon - in roughly forty thousand years. The longest dimension of the complex is only about ten thousand astronomical units (one AU is the average distance of the Earth from the Sun), and so these objects are close enough together for gravity to be the major influence in their development; velocity measurements confirm that the objects are physically associated.

It is possible – even likely - that during the stars' development their orbital motions will prompt the ejection of one or two members from the system, but for now it appears that at least one binary pair will survive for longer times. Other stellar systems need to be examined in

order to see how widespread these young multiplets really are, but the new results support models in which multiple [stars](#) form very early in the stellar womb.

More information: "The formation of a quadruple star system with wide separation" *Nature* 518, 213–215 (12 February 2015) [DOI: 10.1038/nature14166](#)

Provided by Harvard-Smithsonian Center for Astrophysics

Citation: Quadruplets in a stellar womb (2015, February 23) retrieved 22 May 2024 from <https://phys.org/news/2015-02-quadruplets-stellar-womb.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--