

Fewer viral relics may be due to a less bloody evolutionary history

February 2 2015

Humans have fewer remnants of viral DNA in their genes compared to other mammals, a new study has found. This decrease could be because of reduced exposure to blood-borne viruses as humans evolved to use tools rather than biting during violent conflict and the hunting of animals.

Despite natural defence systems, a retrovirus occasionally infects a mammal's egg or sperm, and the virus's genetic code gets incorporated into the animal's own genome. This viral 'fossil' then passes down from generation to generation: we all carry remnants of DNA from viruses that infected our ancestors millions of years ago. These 'endogenous retroviruses' (ERVs) appear not to cause us any harm, even though they are known to result in diseases such as cancer in other animals.

A team of researchers from the University of Oxford and Plymouth University, UK, and the Aaron Diamond AIDS Research Center, USA, wondered if there was a combination of factors unique to humans that explained why these viral fossils in our genomes remain benign. They counted the number of times that retroviruses appear to have been integrated into an animal's genome in humans, comparing humans with 39 other mammalian species, including chimpanzees, dolphins and giant pandas.

Reporting their results in the journal *Retrovirology*, the researchers compared the genetic signature of the two edges of the virus. These edges are identical when the virus first invades the genome, but as they



acquire random mutations over time, they slowly begin to diverge. By tracking this divergence, the research team could measure how long the retrovirus had spent in an animal's genome.

Using this measure, they found that, compared to other animals, far fewer retroviruses were incorporated into the genome for humans and other apes over the last 10 million years. Even compared to animals very similar to us, humans are unusual in not having acquired any new types of retroviruses into their DNA over the last 30 million years.

One reason for the reduction in retroviral incorporation into the <u>human</u> genome might be a change in behaviour as humans evolved: fewer bloody fights and less exposure to infected meat meant that compared to other <u>animals</u>, our ancestors became less likely to encounter blood, a major route for viral infection.

'Considering us simply as a primate species, the proportion of human individuals that are infected with retroviruses is much less than among our relatives such as chimpanzees,' said Dr Robert Belshaw from Plymouth University.

However, lead researcher Dr Gkikas Magiorkinis from Oxford University's Department of Zoology said: 'We have shown in the past that Hepatitis C, a virus transmitted mainly through blood, was spread massively after World War II. There is no doubt that the past trend of reduced blood contacts has been reversed in the last century, and this has severe consequences for viral infections.'

Provided by Oxford University

Citation: Fewer viral relics may be due to a less bloody evolutionary history (2015, February 2) retrieved 7 May 2024 from



https://phys.org/news/2015-02-viral-relics-due-bloody-evolutionary.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.