


Coorong fish hedge their bets for survival

March 26 2015, by Robyn Mills



Sectioned ear bone of black bream showing its growth rings. Credit: Zoe Doubleday, University of Adelaide.

Analysis of the ear bones of the River Murray estuarine fish black bream has revealed how these fish 'hedge their bets' for population survival.

Published in the journal *Biology Letters*, University of Adelaide research has shown that within this single species of [fish](#) there are some

individuals which migrate to different parts of the Coorong in South Australia, and some that generally stay in the one location. Black bream are important for recreational and commercial fishing.

"When we consider animal migration, we tend to think of large seasonal migrations of species like the humpback whale or the Arctic tern. We don't often think of [migratory behaviour](#) that varies within populations," says Professor Bronwyn Gillanders, from the University's School of Biological Sciences and Environment Institute.

"But it appears that within the black bream Coorong population there is a 'bet-hedging strategy' that allows the fish to survive and persist in the Coorong over good times and bad.

"Migration to another area may be more favourable under drought conditions when the water becomes more saline and, conversely, when there is lots of fresh water coming in and there is lots of food readily available, it would be more beneficial for the fish to stay in the location. This probably helps to make the species more resistant to both climate and human-related change."

The researchers used the ear bones of fish collected throughout the estuary to construct their findings. Fish ear bones provide much information through analysis of the trace elements they contain and the width of their growth rings.

"Like tree growth rings, the [ear bones](#) reveal the age of the fish and growth periods which correlate with the growth of the fish itself," says Professor Gillanders. "When we measure the width of the growth increments, we can trace back to see how fast the fish was growing at a particular time and year.

"The bones can also tell us whether the fish is migratory or 'resident' by

mapping the ratios of barium against calcium. The higher levels of barium indicate when the fish was in fresher water."

Professor Gillanders found that 62% of the fish were resident and 38% were migratory. Models were used to investigate differences in annual growth between the two groups and construct a growth time series.

"Throughout the late 1990s and early 2000s resident fish had increased growth compared with migrant fish but this changed around 2005 when growth of migrant fish increases," says Professor Gillanders. "This is likely to be a result of the deteriorating conditions in the Coorong and reflects the ability of the migratory fish to find more favourable conditions and source more food."

More information: "Partial migration: growth varies between resident and migratory fish." [DOI: 10.1098/rsbl.2014.0850](https://doi.org/10.1098/rsbl.2014.0850)

Provided by University of Adelaide

Citation: Coorong fish hedge their bets for survival (2015, March 26) retrieved 26 April 2024 from <https://phys.org/news/2015-03-coorong-fish-hedge-survival.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.