

More evidence for groundwater on Mars

March 27 2015



Figure 1 is from Pondrelli et al. (A) Location map of the study area on MOLAbased shaded relief map. Topographic contours (in black, 1000 m spacing) are indicated. (B) High Resolution Stereo Camera (HRSC) mosaic of the mapped area. Topographic contours (in white, 500 m spacing) are indicated. (C) Excerpt of the geological map by Scott and Tanaka (1986) on an HRSC mosaic. Geologic units (see text for more details): Npl1--Noachian cratered unit of the plateau sequence; Npl2--Noachian subdued crater unit of the plateau sequence; Hr--Hesperian ridged plains material. (D) Footprints of High Resolution Imaging Science Experiment (HiRISE) coverage on an HRSC mosaic. The white fi lling indicates stereo pairs. The area is fully covered by Context Camera (CTX) imagery. Compact Reconnaissance Imaging Spectrometer for Mars (CRISM) scenes used in this study are recognizable by the hourglass shape. Credit: Pondrelli et al. and GSA Bulletin



Monica Pondrelli and colleagues investigated the Equatorial Layered Deposits (ELDs) of Arabia Terra in Firsoff crater area, Mars, to understand their formation and potential habitability. On the plateau, ELDs consist of rare mounds, flat-lying deposits, and cross-bedded dune fields. Pondrelli and colleagues interpret the mounds as smaller spring deposits, the flat-lying deposits as playa, and the cross-bedded dune fields as aeolian. They write that groundwater fluctuations appear to be the major factor controlling ELD deposition.

Pondrelli and colleagues also note that the ELDs inside the craters would likely have originated by fluid upwelling through the fissure ridges and the <u>mounds</u>, and that lead to evaporite precipitation. The presence of spring and playa deposits points to the possible presence of a hydrological cycle, driving groundwater upwelling on Mars at surface temperatures above freezing. Pondrelli and colleagues write that such conditions in a similar Earth environment would have been conducive for microbial colonization.

As a basis for their research, Pondrelli and <u>colleagues</u> produced a detailed geological map of the Firsoff crater area. The new map includes <u>crater</u> count dating, a survey of the stratigraphic relations, and analysis of the depositional geometries and compositional constraints. They note that this ELD unit consists of sulfates and shows other characteristics typical of evaporites such as polygonal pattern and indications of dissolution.

More information: Equatorial layered deposits in Arabia Terra, Mars: Facies and process variability, M. Pondrelli et al., International Research School of Planetary Sciences, Università d'Annunzio, Pescara, Italy. Published online ahead of print on 10 Mar. 2015; <u>http://dx.doi.org/10.1130/B31225.1</u>.



Provided by Geological Society of America

Citation: More evidence for groundwater on Mars (2015, March 27) retrieved 2 May 2024 from <u>https://phys.org/news/2015-03-evidence-groundwater-mars.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.