

Researchers fine-tune quantum dots from coal

March 18 2015



Vials hold solutions with graphene quantum dots that fluoresce in different colors depending on the dots' size. Techniques to produce the dots in specific sizes using coal as a source were developed at Rice University. Credit: Tour Group/Rice University

Graphene quantum dots made from coal, introduced in 2013 by the Rice University lab of chemist James Tour, can be engineered for specific semiconducting properties in either of two single-step processes.

In a new study this week in the American Chemical Society journal *Applied Materials & Interfaces*, Tour and colleagues demonstrated fine control over the graphene oxide dots' size-dependent band gap, the property that makes them semiconductors. Quantum dots are semiconducting materials that are small enough to exhibit quantum



mechanical properties that only appear at the nanoscale.

Tour's group found they could produce quantum dots with specific semiconducting properties by sorting them through ultrafiltration, a method commonly used in municipal and industrial water filtration and in food production.

The other single-step process involved direct control of the reaction temperature in the oxidation process that reduced coal to quantum dots. The researchers found hotter temperatures produced smaller dots, which had different semiconducting properties.

Tour said graphene quantum dots may prove highly efficient in applications ranging from medical imaging to additions to fabrics and upholstery for brighter and longer-lasting colors. "Quantum dots generally cost about \$1 million per kilogram and we can now make them in an inexpensive reaction between coal and acid, followed by separation. And the coal is less than \$100 per ton."

The dots in these experiments all come from treatment of anthracite, a kind of coal. The processes produce batches in specific sizes between 4.5 and 70 nanometers in diameter.





Rice University scientists have produced graphene quantum dots produced from coal with tuned band gaps and photoluminescent properties. These quantum dots, seen with an electron microscope, average 70 nanometers in diameter. Credit: Tour Group/Rice University

Graphene quantum dots are photoluminescent, which means they emit light of a particular wavelength in response to incoming light of a different wavelength. The emitted light ranges from green (smaller dots) to orange-red (larger dots). Because the emitted color also depends on the dots' size, this property can also be tuned, Tour said. The lab found



quantum dots that emit blue light were easiest to produce from bituminous <u>coal</u>.

The researchers suggested their quantum dots may also enhance sensing, electronic and photovoltaic applications. For instance, catalytic reactions could be enhanced by manipulating the reactive edges of <u>quantum dots</u>. Their fluorescence could make them suitable for metal or chemical detection applications by tuning to avoid interference with the target materials' emissions.



Rice University scientists have produced graphene quantum dots produced from



coal with tuned band gaps and photoluminescent properties. These quantum dots are about 4.5 nanometers in diameter. Credit: Tour Group/Rice University

More information: *Applied Materials & Interfaces*, <u>pubs.acs.org/doi/abs/10.1021/acsami.5b01419</u>

Provided by Rice University

Citation: Researchers fine-tune quantum dots from coal (2015, March 18) retrieved 2 May 2024 from <u>https://phys.org/news/2015-03-fine-tune-quantum-dots-coal.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.