


# Measuring galaxy evolution with globular clusters

March 23 2015

---



The galaxy NGC 4365 in the Virgo supercluster of galaxies, showing its many globular clusters (most of the more than one hundred dots in this image are actually globular star clusters). A new study of over seven thousand globular

clusters around ten Virgo galaxies finds that they are often gathered into distinct groupings whose shapes reflect the formation of the clusters and the histories of the galaxies. Credit: NASA/HST and ESO VLT

Globular clusters are gravitationally bound ensembles of stars, as many as a million stars in some cases, grouped in roughly spherical clusters with diameters as small as only tens of light-years. Globular clusters are typically located in the outer regions (the halos) of galaxies; the Milky Way galaxy has about two hundred globular clusters orbiting it. Astronomers are interested in globular clusters in part because they are home to many of the oldest known stars, but also because of their locations in the halos.

Collisions between [galaxies](#) are commonplace, and globular clusters may provide fossil evidence of these encounters because they are strongly affected by such interactions. During a collision, a galaxy can grow by absorbing or merging with its neighbor, and some models predict that clusters form during these interactions. Moreover, it is possible that in a merger large numbers of globular clusters originally belonging to a smaller galaxy may be captured by the larger galaxy. In any case, the distribution of globular clusters around a galaxy holds clues to their origins and the history of its host galaxy.

The Virgo Cluster of galaxies, containing between one and two thousand galaxies, is located about fifty-four million light-years away in the direction of the constellation of Virgo. The ten brightest galaxies of the Virgo Cluster alone contain 7053 detected globular clusters. CfA astronomers Raffaele D'Abrusco, Pepi Fabbiano, and Andreas Zezas carefully examined this set of globular clusters looking for information about the history of these galaxies. In a new paper, they report discovering distinctive structures among the globular cluster systems,

meaning that the [globular clusters](#) around these galaxies are not distributed symmetrically. Their configurations often take shapes ranging from roughly linear to circular, with some more complex shapes as well. The scientists found 229 such structures in this subsample, forty-two of them classified as being medium or large and stretching over as much as seventy-five light-years. The elongated structures tend to be aligned with an axis of the [host galaxy](#), as would be expected if a merger were responsible.

The scientists argue that these structures are indeed the remnants of galaxies that were accreted in the past, and among other things they estimate limits on the masses of these [parent galaxies](#). Computer simulations provide some rough level of agreement. The authors note that with more detailed computations, these structures offer a powerful new tool to advance the study of [galaxy evolution](#).

**More information:** "Spatial Structures in the Globular Cluster Distribution of the Ten Brightest Virgo Galaxies," R. D'Abrusco, G. Fabbiano, A. Zezas, *ApJ* 2015 (in press) [arxiv.org/abs/1503.04819](https://arxiv.org/abs/1503.04819)

Provided by Harvard-Smithsonian Center for Astrophysics

Citation: Measuring galaxy evolution with globular clusters (2015, March 23) retrieved 18 April 2024 from <https://phys.org/news/2015-03-galaxy-evolution-globular-clusters.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--