


# Methane oxidation fuelled by algal oxygen production

March 3 2015

---



Sampling was performed from a platform on Lake Cadagno, Switzerland

Methane emissions are strongly reduced in lakes with anoxic bottom waters. But here – contrary to what has previously been assumed – methane removal is not due to archaea or anaerobic bacteria. A new study on Lake Cadagno in Canton Ticino shows that the microorganisms responsible are aerobic proteobacteria. The oxygen they require is

produced in situ by photosynthetic algae.

In contrast to oceans, [freshwater lakes](#) – and tropical reservoirs – are significant sources of [methane](#) emissions. Methane, a greenhouse gas, arises from the degradation of organic material settling on the bottom. Although lakes occupy a much smaller proportion of the Earth's surface than oceans, they account for a much larger proportion of [methane emissions](#). Well-mixed lakes, in turn, are the main contributors, while emissions from seasonally or permanently stratified lakes with anoxic bottom waters are greatly reduced. It has been assumed to date that the methane-removing processes occurring in such lakes are the same as those in marine systems. But a new study carried out on Lake Cadagno (Canton Ticino) by researchers from Eawag and the Max Planck Institute for Marine Microbiology (Bremen, Germany) shows that this is not the case.

The scientists demonstrated that methane is almost completely consumed in the anoxic waters of Lake Cadagno, but they did not detect any known anaerobic methane-oxidizing [bacteria](#) – or archaea, which are responsible for marine methane oxidation. Instead, water samples collected from a depth of around 12 metres were found to contain abundant aerobic proteobacteria – up to 240,000 cells per millilitre.

"We wondered, of course, how these aerobic bacteria can survive in anoxic waters," says first author Jana Milucka of the Max Planck Institute for Marine Microbiology. To answer this question, the behaviour of the bacteria was investigated in laboratory experiments: methane oxidation was found to be stimulated only when oxygen was added to the samples incubated in vitro, or when they were exposed to light. The scientists concluded that the oxygen required by the bacteria is produced by photosynthesis in neighbouring diatoms. Analysis by fluorescence microscopy showed that methane-oxidizing bacteria belonging to the family Methylococcaceae occur in close proximity to

diatoms and can thus utilize the oxygen they generate.

Thanks to the combined activity of bacteria and diatoms, methane is thus consumed in the [lake](#) rather than being released into the atmosphere. This type of methane removal has not previously been described in freshwater systems. Project leader Carsten Schubert of Eawag comments: "For lakes with anoxic layers, and also for certain marine zones, it looks as if the textbooks will have to be rewritten." Aerobic methane-oxidizing bacteria may play a significant role wherever sufficient light penetrates to anoxic water layers; according to Schubert, this is the case in most Swiss lakes. Similar observations have already been made in Lake Rotsee near Lucerne, in studies not yet published. Research will now focus on deeper lakes, where initial investigations suggest that different processes occur.

**More information:** "Methane oxidation coupled to oxygenic photosynthesis in anoxic waters." *The ISME Journal* , (13 February 2015) | [DOI: 10.1038/ismej.2015.12](https://doi.org/10.1038/ismej.2015.12)

Provided by Swiss Federal Institute of Aquatic Science and Technology

Citation: Methane oxidation fuelled by algal oxygen production (2015, March 3) retrieved 18 April 2024 from <https://phys.org/news/2015-03-methane-oxidation-fuelled-algal-oxygen.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--