

Spurring production of a sluggish enzyme for crop yields

March 3 2015



Maxim Kapralov (L) and Spencer Whitney. Credit: Charles Tambiah / ARC-CoETP

Australian scientists have found a way to improve production of an enzyme essential to plant growth.

The discovery advances efforts to improve global food security that aim to increase the yields of some of our most important staple crops, such as wheat, cotton and rice.

"The enzyme Rubisco has baffled scientists for over 50 years, as it is thousands of times less efficient at its job than most other enzymes," said Dr Spencer Whitney from The Australian National University (ANU) who was leader of the research team from the ARC Centre of Excellence for Translational Photosynthesis.

Much of the confusion stems from the highly complex structure of Rubisco. It is an enzyme with 16 pieces that requires the assistance of more than 12 other proteins to assemble it correctly.

"We focussed on one of Rubisco's partner proteins, called RAF1. By supplying a Rubisco with its partnering RAF1 we found the amount of Rubisco produced in leaves could be more than doubled," he said.

Rubisco is the most abundant protein on the planet. It also has one of the most significant responsibilities in life on Earth: the conversion of carbon dioxide to organic compounds - the building blocks of all plant and animal cells.

"Understanding the partnership with RAF1 has important implications with regard to ongoing efforts to accelerate the sluggish activity of Rubisco - a key goal for improving crop productivity," Dr Whitney said.

"In a world with increasing demands for food, this is a milestone towards increasing the photosynthetic rate in crop plants that rely heavily on Rubisco."

The study, done in collaboration with the University of Wollongong and published in the latest *PNAS* journal, shows Rubisco and RAF1 have evolved together. Changes in one protein requires a balancing change in the other.

"We improved the production of a modified Rubisco in leaves by

inserting a modified complimentary version of RAF1," Dr Whitney said.

"Including RAF1 caused Rubisco's levels to double, and as a result, we obtained faster rates of photosynthesis and [plant growth](#) compared to the plants where RAF1 was not introduced.

"The finding also explains why our prior attempts to insert more efficient versions of Rubisco from some algae into [plant leaves](#) have failed – they require different chaperones to those available in leaf cells."

More information: Improving recombinant Rubisco biogenesis, plant photosynthesis and growth by coexpressing its ancillary RAF1 chaperone, *PNAS*, www.pnas.org/cgi/doi/10.1073/pnas.1420536112

Provided by Australian National University

Citation: Spurring production of a sluggish enzyme for crop yields (2015, March 3) retrieved 2 May 2024 from <https://phys.org/news/2015-03-spurring-production-sluggish-enzyme-crop.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--