

Preventing Australia's biodiversity from collapsing

April 8 2015



eadbeater's possum. Credit: Takver on Flickr

Preventing Australia's biodiversity from collapsing needs long-term science-based programs that are more complex than just culling cats and foxes, says leading environmental scientist Professor David Lindenmayer.

Professor Lindenmayer from The Australian National University (ANU)

said short-term programs could sometimes be ineffective or even detrimental.

"You have to match up the problem to the solution to find the most effective approach," said Professor Lindenmayer, from the ANU Fenner School of Environment and Society.

"Otherwise you're just going to waste a whole lot of [public money](#)."

Professor Lindenmayer's views have been published as a commentary in the latest *PNAS* journal.

Australian mammals have gone extinct in the last 200 years at 10 times the rate of those in the US, with 20 per cent of the mammals remaining now under threat.

Other species survive, but only in a tiny fraction of their previous range, said Professor Lindenmayer.

"The bilby is associated with arid areas these days, but it used to flourish in woodlands such as those around Canberra," he said.

Feral cats and foxes are a problem in many parts of Australia. However, locations throughout the country face other key threats.

"The Leadbeater's possum in Victoria is at a real risk of global extinction as a result of industrial clearfelling and recurrent [wildfires](#) that are exacerbated by past logging history," Professor Lindenmayer said.

"The science is there in many cases. For many animals we know what to do.

"For example large old trees are keystone structures in forests and

savannas, more than 300 species cannot survive without them," he said.

Australia leads the world in extinctions, but could lead the world in conservation, due to its wealth, its continent-wide government and the quality of its scientists.

"We urgently need to do more targeted management to prevent declines and monitoring with trigger points for action when species go into decline," Professor Lindenmayer said.

"We also need monitoring so we can stop the next generation of feral invaders, for example beasts brought in through the pet trade."

More information: "Continental-level biodiversity collapse." *PNAS* 2015 ; published ahead of print April 6, 2015, [DOI: 10.1073/pnas.1502766112](https://doi.org/10.1073/pnas.1502766112)

Provided by Australian National University

Citation: Preventing Australia's biodiversity from collapsing (2015, April 8) retrieved 17 April 2024 from <https://phys.org/news/2015-04-australia-biodiversity-collapsing.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.