

# Grey squirrels' role as hosts of Lyme disease bacteria under the spotlight

April 22 2015, by Stuart Forsyth

---



Credit: Aileen Adam, University of Glasgow

Grey squirrels have been described as one of the 'world's worst invasive species' and have caused a decline in indigenous red squirrel populations and damaged forestry in the UK.

Now the role of this [invasive species](#) in hosting the bacteria which causes Lyme disease in humans has come under the spotlight in a study by ecologists at the University of Glasgow.

In Britain, Lyme disease is caused by four [species](#) of the bacteria *Borrelia burgdorferi sensu lato*. These bacteria are maintained in a tick-wildlife cycle and a bite from an infected tick is responsible for Lyme disease in people.

In the UK, the main tick species which transmits Lyme disease, *Ixodes ricinus*, is commonly known as the deer or sheep tick. Around 4% of ticks in the UK are infected, though this can vary substantially between different areas. Species of *Borrelia* bacteria can infect a number of wildlife species, but generally birds and rodents are considered to be the most important hosts, while deer are important for maintaining tick populations but do not become infected with the bacteria.

Caroline Millins, of the Institute of Biodiversity, Animal Health and Comparative Medicine and lead author of the study published in *Applied and Environmental Microbiology*, a journal published by the American Society for Microbiology, said: "Grey squirrels are abundant and widely distributed in the UK, so we were interested in finding out how this invasive species may contribute to maintaining the bacteria which causes Lyme disease in humans."

Grey squirrels could act as an alternative host for ticks. Based on other studies of invasive species, they may either increase or decrease the number of infected ticks in the environment, an important risk factor for human infections.

The researchers tested 679 squirrels from across Scotland and the North of England in 2012 and 2013. They found that [grey squirrels](#) often carried ticks, and could be infected with all four of the *Borrelia* species found in the UK. Around 12% of the collected squirrels were infected, and surprisingly, the most frequent infection was a species usually found in birds.

Miss Millins said: "Frequent infection of grey squirrels with bird strains of *Borrelia* was unexpected, and challenges our current understanding of host pathogen interactions for this zoonotic pathogen.

"We found quite different patterns and duration of infection in grey squirrels compared to native woodland rodent species. Grey squirrels become infected with whichever strains are circulating in the local area, and our models suggest that the duration of infection isn't life-long. In contrast, native rodents tend to develop chronic lifelong infections with strains of *Borrelia* that have adapted to these hosts.

"We can't say from this study whether grey squirrels lead to an increase or decrease in the number of ticks infected with *Borrelia* species in an area, but we have produced comprehensive baseline data that future studies could use to investigate this."

Grey [squirrels](#) were introduced to the UK between 1876 and 1929 and are regarded as a pest species under national legislation.

Lyme disease was first diagnosed in Old Lyme, Connecticut, in the US, in 1975. The most common symptom is a distinctive skin rash which has a 'target' or 'bull's eye' appearance. Patients can also have flu-like symptoms, including a headache and muscle pains. Symptoms can start within 3–30 days of an infected tick [bite](#).

If early symptoms of Lyme disease aren't recognised and treated, infection can spread to cause more serious systemic chronic inflammatory disease, often affecting the joints, nervous system or heart. Lyme disease is treated with antibiotics.

The best way to prevent Lyme disease is to check yourself for ticks as soon as possible after walking or cycling in known tick areas. Ticks should be removed carefully with tweezers, or ideally with a tick

remover available from most veterinary clinics.

**More information:** For more information on tick prevention go to Borreliosis and Associated Diseases Awareness UK, [www.bada-uk.org/](http://www.bada-uk.org/)

Provided by University of Glasgow

Citation: Grey squirrels' role as hosts of Lyme disease bacteria under the spotlight (2015, April 22) retrieved 19 April 2024 from <https://phys.org/news/2015-04-grey-squirrels-role-hosts-lyme.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.