

# High mountains warming faster than expected

April 23 2015

---



Elevation-dependent warming is a poorly observed phenomenon that requires urgent attention to ensure that potentially important changes in high mountain environments are adequately monitored by the global observational network," say members of the Mountain Research Initiative Working Group. Credit: Douglas Hardy, UMass Amherst

High elevation environments around the world may be warming much faster than previously thought, according to members of an international research team including Raymond Bradley, director of the Climate System Research Center at the University of Massachusetts Amherst. They call for more aggressive monitoring of temperature changes in mountain regions and more attention to the potential consequences of warming.

"Elevation-dependent [warming](#) is a poorly observed phenomenon that requires urgent attention to ensure that potentially important changes in high mountain environments are adequately monitored by the global observational network," say members of the Mountain Research Initiative Working Group in the current issue of *Nature Climate Change*.

High mountains are the major water source for large numbers of people living at lower elevations, so the social and [economic consequences](#) of enhanced warming in mountain regions could be large, the researchers add. "This alone requires that close attention be paid to the issue. In addition, mountains provide habitat for many of the world's rare and endangered species, and the presence of many different ecosystems in close proximity enhances the ecological sensitivity of mountains to environmental change."

Lead author Nick Pepin of the University of Portsmouth, U.K., says, "There is growing evidence that high mountain regions are warming faster than lower elevations and such warming can accelerate many other environmental changes such as glacial melt and vegetation change, but scientists urgently need more and better data to confirm this. If we are right and mountains are warming more rapidly than other environments, the social and economic consequences could be serious, and we could see more dramatic changes much sooner than previously thought."

UMass Amherst's Bradley adds that without substantially better

information, there is a risk of underestimating the severity of a number of problems, including water shortages and the possible extinction of some alpine flora and fauna.

He says, "We are calling for special efforts to be made to extend scientific observations upwards to the highest summits to capture what is happening across the world's mountains. We also need a strong effort to find, collate and evaluate observational data that already exists wherever it is in the world. This requires international collaboration."

Records of weather patterns at high altitudes are "extremely sparse," the researchers found. There are very few weather stations above 14,700 feet (4,500 m), and long-term data, crucial for detecting patterns, doesn't yet exist above 16,400 feet (5,000 m) anywhere in the world. The authors say the longest observations above this elevation are from the summit of Kilimanjaro, which have been maintained for more than a decade by Douglas Hardy of UMass Amherst.

For this study, Pepin, Bradley and colleagues reviewed elevation-dependent warming mechanisms such as loss of snow and ice, increased latent heat release at high altitudes, low-elevation aerosol pollutants that increase the difference in warming rates between low and high elevations, plus other factors that enhance warming with elevation in different regions, and in different seasons.

They discuss future needs to improve knowledge of mountain temperature trends and mechanisms via improved observations, satellite-based remote sensing and model simulations. Noting that "many factors make it extremely difficult to determine the rate of warming in mountainous regions," the team reports the most striking evidence that [mountain regions](#) are warming more rapidly than surrounding regions comes from the Tibetan plateau, where temperatures have risen steadily over the past 50 years and the rate of change is accelerating.

This research team with members from the U.K., U.S., Switzerland, Canada, Ecuador, Pakistan, China, Italy, Austria and Kazakhstan, came together as part of the Mountain Research Initiative, a mountain global change research effort funded by the Swiss National Foundation.

**More information:** Elevation-dependent warming in mountain regions of the world, *Nature Climate Change* 5, 424–430 (2015) [DOI: 10.1038/nclimate2563](https://doi.org/10.1038/nclimate2563)

Provided by University of Massachusetts Amherst

Citation: High mountains warming faster than expected (2015, April 23) retrieved 20 April 2024 from <https://phys.org/news/2015-04-high-mountains-faster.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--