


All will be illuminated: Real-time multicolor imaging with luminescent protein-based Nano-lanterns

April 22 2015, by Stuart Mason Dambrot



Luminescence imaging of intracellular microstructures. (A) Luminescence and fluorescence images of subcellular structures: nuclei in interphase (CNL-H2B and ONL-H2B), F-actin [inositol trisphosphate 3-kinase A (ITPKA) -ONL], microtubules (β -tubulin-ONL), peroxisomes [peroxisome targeting signal 2 (PTS2) -CNL], and focal adhesions (zyxin-ONL). (B) Subnuclear structures

visualized by dual-color luminescence imaging: CNL-fibrillarin (nucleoli) and ONL-H2B. Each luminescence signal was separated by linear unmixing. (Scale bars: 10 μm .) Credit: Takai A, et al (2015) Expanded palette of Nano-lanterns for real-time multicolor luminescence imaging. *Proc Natl Acad Sci USA* 112(14):4352-4356.

While fluorescence imaging (in which external light is used to excite a specimen that then emits light in response) is essential in cell biology, it has a number of significant drawbacks, including autofluorescence, phototoxicity and photobleaching, resulting from that excitation light. In addition, fluorescence imaging has the unfortunate side effect of triggering cellular activation when combined with optogenetics – an otherwise extremely valuable tool. On the other hand, [luminescence](#) (in this case, a type of chemiluminescence called bioluminescence) imaging doesn't require light activation, and so eschews these issues – but currently suffers from low brightness and poor color variants.

Recently, however, scientists at RIKEN and Osaka University extended their previous development of a bright yellowish-green luminescent protein *Nano-lantern* to devise bright cyan and orange luminescent proteins some 20 times brighter than previously possible with wild-type (i.e., naturally-occurring) *Renilla luciferase*, or *Rluc* – an oxidative enzyme associated with a luciferin-binding protein – found in a type of soft coral known as a sea pansy. (*Luciferins* are organic substances, found in luminescent organisms, which produce a near-heatless light upon oxidation.) Specifically, the researchers accomplished this by bioluminescence resonance energy transfer (BRET) from enhanced *Renilla luciferase* to a fluorescent protein, stating that their proof-of-principle experiments show that luminescence imaging has become a practical alternative when the side effects by the excitation light are not negligible – for example, when the samples are very sensitive to

photodamage – and that the most effective future application of luminescence imaging lies in combining it with optogenetics, since the latter's external light illumination can be reserved for optical stimulation.

Dr. Yasushi Okada discussed the paper that he, Dr. Akira Takai, Dr. Takeharu Nagai, and their colleagues published in *Proceedings of the National Academy of Sciences*, starting with the challenges of developing cyan and orange luminescent proteins approximately 20 times brighter than wild-type *Renilla luciferase*. "Making the cyan version was straightforward," Okada tells *Phys.org*. "We acquired the best available cyan fluorescent protein, *mTurquoise2*," or *mTq2*, "from Dr Joachim Goedhart at University of Amsterdam. I was so impressed with the title of his paper¹ that I immediately requested the plasmid – and he said that my request was the first he received. Several weeks after, we started working with our colleague Dr. Takeharu Nagai on yellow lanterns and soon came up with the idea of swapping the *mVenus* used in the yellow Nano-lantern," or YNL, "with *mTq2* – and it worked excellently, producing the cyan Nano-lantern," or CNL.

That said, Okada adds that developing the orange Nano-lantern (ONL) took trial-and-error. "We initially planned to use a large Stokes shift orange fluorescent protein like *LSSmOrange2*, but it didn't work well." A Stokes shift, which is essential in [fluorescence imaging](#), refers to the difference between the energy (i.e., wavelength) of the excitation and emitted photons – and a large Stokes shift typically indicates a greater difference and thereby easier detection. "We therefore tested all possible combinations of available orange to red fluorescent proteins and *Rluc* variants – hundreds of them – and we finally identified the best combination." That combination became the orange Nano-lantern.


The paper describes an application in which three gene expression reporters (genes attached to a regulatory sequence of a gene being investigated) were introduced into the genome of photosensitive

embryonic stem (ES) cells by using *transposase mediated insertion*. (*Transposase* is an enzyme that binds to the end of a transposon – a DNA sequence that can change its position within the genome – and catalyzes transposon movement to and insertion in another part of the genome.) The triple positive cells were then screened by fluorescence-activated cell sorting (FACS), a method for sorting a heterogeneous mixture of biological cells into two or more containers, one cell at a time, based upon the specific light scattering and fluorescent characteristics of each cell. "Here, the dual use of our Nano-lanterns was very useful," Okada recounts. "They can be observed both by fluorescence and luminescence, so we can therefore use FACS for screening for the triple positive cells. "Our major imaging challenge was the consumption of the substrate, coelenterazine." The light detected in bioluminescence imaging is produced when luciferase is expressed *in vivo* and oxidizes a molecular substrate, while coelenterazine is the luciferin found in many aquatic organisms.

"Although we've developed diacetyl coelenterazine," in which protective acetyl groups inhibit autoxidation, allowing it to be added at higher concentrations without increasing background autoluminescence, "we needed to supply it continuously for long-term observation that exceed 12 hours. Our colleague and co-author Dr. Tomonobu Watanabe performed the imaging with his handmade perfusion system."

Another issue in luminescence imaging with Nano-lanterns is that its signal intensity is still over 100 times weaker than that of fluorescent proteins. "Yes, our Nano-lanterns are still darker than fluorescent proteins," Okada agrees, "but fluorescent protein brightness is proportional to the excitation power until saturation. With strong excitation close to saturation, a single fluorescent protein molecule can emit more than 10^7 photons per second, while our Nano-lantern emits less than 10 photons per second. However, to avoid photobleaching we normally do not use such strong excitation beams for fluorescent live cell

imaging. Even for a single molecule live cell imaging" he points out, "we reduce the excitation power to get only 100-1000 photons per second from a single [fluorescent protein](#). In short, the "more than 100 times weaker" comparison in the article is based on this calculation – and considering that most of live cell fluorescent imaging is normally done with much weaker excitation beams, Nano-lantern brightness in such applications will be much closer to that of fluorescent proteins."



Inhomogeneous expression of pluripotency markers in a single colony of ES cells. Luminescence signals of reporters for (A) Oct4 (CNL), (B) Nanog (YNL), and (C) Sox2 (ONL) were separated by linear unmixing and overlaid with the bright-field image. (D) Signals of CNL, YNL, and ONL were merged and overlaid with the bright-field image. (Scale bars: 100 μm .) Credit: Takai A, et al (2015) Expanded palette of Nano-lanterns for real-time multicolor luminescence imaging. *Proc Natl Acad Sci USA* 112(14):4352-4356.

Okada acknowledges that since luminescence imaging requires a molecular substrate instead of excitation light, and the supply and consumption of the substrate limit imaging time, further increases in luminescence imaging light output would require higher enzymatic turnover rate because the number of photons per single enzymatic reaction does not exceed 1. The substrate would then be exhausted more rapidly, making substrate supply the more important challenge. The next frontier is therefore enhancing substrate synthesis or recycling. "For the bacterial luciferase and firefly luciferase" (the former is found in the *Photobacterium* species *Vibrio fischeri*, *V. haweyi*, and *V. harveyi*) "the supply systems for their substrate are already reconstituted in bacteria or plants. It won't take much time before the coelenterazine regeneration system will be reconstituted *in vitro* or *in vivo*."

Luminescence imaging also lacks optical sectioning capabilities. "If you use a conventional microscope," Okada explains, "the axial resolution of luminescence imaging is much poorer than fluorescence imaging. (Axial resolution refers to the ability to image at various depths in 3D imaging.) "However, we have several ideas to increase axial resolution, because it's critically important for applications like *in vivo* imaging."

In fluorescent imaging, different colors of fluorescent proteins can easily be separated by the combination of excitation and emission filters,

whereas in luminescence imaging signals from different colors of Nano-lanterns can be separated only by the emission spectrum, and therefore require (albeit a simpler) linear unmixing algorithm. "While the emission spectra of luminescent proteins were much broader than fluorescent proteins, Nano-lantern emission is essentially same as that of the fluorescent proteins," Okada tells *Phys.org*. "As a result, the image is sharp enough for efficient and accurate separation by the linear unmixing algorithm. Furthermore, the background is completely dark, because luminescence imaging is free from autofluorescence, making the luminescence images are even more suitable for post-processing unmixing." (*Autofluorescence* is the natural emission of light by biological structures after they absorb light, and is used to distinguish the light originating from artificially added fluorescent markers called fluorophores.) "Related to this background issue," relates Okada, "we received a query from our paper's *PNAS* editor that the images were apparently manipulated because the background is homogeneously dark. We pointed out that this extremely low level background is a feature of luminescence imaging that makes it more sensitive and quantitative."

Despite the challenges they met and still face, Okada says that "From our experiences in fluorescence imaging, as well as the history of the development of fluorescent proteins, it was clear from the beginning that development of new color variants was absolutely required to extend the application of luminescence imaging – and fortunately, the production methodologies for color variants were evident from the BRET-based design of the yellow Nano-lantern."

It is also the case that Nano-lantern luminescence imaging has a range of significant benefits relative to current luminescence imaging, one being its ability to visualize rapid dynamics of endosomes and peroxisomes at a temporal resolution of one second. (An *endosome* is a membrane-bounded compartment inside eukaryotic cells involved in endocytic membrane transport, a pathway by which by which cells absorb

molecules by engulfing them; a *peroxisome* is a small, membrane-enclosed organelle containing enzymes involved in a variety of metabolic reactions.) "Previous luminescence probes were very dim, and their imaging normally required extremely long exposure time – typically 10 minutes or longer," Okada points out. "Therefore, extremely low noise cameras, such as those cooled with liquid nitrogen, were required. Alternatively, you have to compromise spatial resolution to increase the frame rate. Therefore, it was impossible to observe small, but rapidly moving structures with luminescence. However, when expressed in animal cells our Nano-lantern is much brighter by several orders of magnitude." In fact, Okada relates, when his collaborators first used Nano-lanterns, they often reported that they could not take images – but this occurred because that their imaging system was optimized for conventional luminescence probes, and so was oversaturated by the Nano-lantern's brightness. "I often advised them to image the sample as if it were fluorescently stained, but without excitation light. That solved the problem."

The endosome and peroxisome experiments were in essence designed to demonstrate Nano-lantern brightness. Since both endosomes and peroxisomes are small intracellular structures with

Citation: All will be illuminated: Real-time multicolor imaging with luminescent protein-based Nano-lanterns (2015, April 22) retrieved 19 July 2024 from <https://phys.org/news/2015-04-illuminated-real-time-multicolor-imaging-luminescent.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.