

## Birds time breeding to hit 'peak caterpillar'

May 25 2015, by Pete Wilton, Oxford Science Blog

---



When oaks burst into life in spring populations of oak-leaf-eating caterpillars boom: this offers a food bonanza for caterpillar-munching birds looking to raise a family.

But, if you're a bird, how do you time your breeding to exploit this seasonal food source? Understanding this is essential if scientists are to predict which species will be able to adapt to spring arriving earlier due to climate change.

Amy Hinks, Ella Cole, and Ben Sheldon, of Oxford University's Department of Zoology, looked into this question by studying great tits at Wytham Woods which feed on winter moth [caterpillars](#) which, in turn, feed on the newly-emerged leaves of oak trees.

The team monitored the bud development of all [oak trees](#) and the timing of egg laying of all great tits across 28-hectares of deciduous woodland. They also measured caterpillar abundance at a sample of oaks throughout spring.

The team report in *American Naturalist* that the timing of leaf emergence of a given oak is a reliable predictor of when caterpillars are most abundant on its foliage. However, individual oaks have their own sense of timing: when each tree buds is consistent over many years and does not tend to follow average temperature trends.

'We found that the laying date is best predicted by the timing of oak leaf emergence within the immediate vicinity, less than a 50 metre radius, of the nest,' said report author Ella Cole of Oxford's Edward Grey Institute.

'There is evidence to suggest that great tits use large-scale cues such as temperature and day length to time their breeding. The results from this study show that temperature cannot be the whole story: in addition to using these global cues, tits are able to fine-tune their breeding decisions based on information from their local environment.

'It is thought that female tits may be using information on the presence of early-stage caterpillars or the stage of development of tree buds. In early spring tits can be seen closely inspecting, and sometimes even eating, tree buds which may be how they collect this information.'

Although there is evidence from other bird species that vegetation cues are used to time egg laying, experiments on captive [great tits](#) that have

tried to test this – by providing birds with either leafing or non-leafing branches – have been unable to influence laying date.

'However, as these experiments were done in captivity, it is not clear how much they reflect what birds do in the wild,' Ella tells me. 'Precisely what cues birds are using to synchronise their breeding cycle with their immediate environment remains a bit of a mystery, and we clearly need further work exploring this question.'

**More information:** "Scale-dependent phenological synchrony between songbirds and their caterpillar food source." *The American Naturalist*, online in advance of print. [dx.doi.org/10.1086/681572](https://doi.org/10.1086/681572)

Provided by Oxford University

Citation: Birds time breeding to hit 'peak caterpillar' (2015, May 25) retrieved 19 April 2024 from <https://phys.org/news/2015-05-birds-peak-caterpillar.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.