


# Chameleon proteins make individual cells visible

May 19 2015

---



Motor neurons of a living zebrafish larva in the spinal cord. One cell, including its body and extensions, is visible – in stark contrast to its green neighbouring cells – in magenta and white. Credit: Microscopic image: Dempsey et al. Nature Methods 2015

Researchers discovered a new mechanism of how fluorescent proteins can change colour. It enables the microscopic visualization of individual cells in their three-dimensional environment in living organisms.

Researchers at ETH Zurich's Department of Biosystems Science and Engineering in Basel have developed a new microscopy technique that enables for the first time to selectively visualize [individual cells](#) within the complex, three-dimensional tissue of a living organism. The researchers have thus succeeded in capturing spectacular microscopic images, such as in the nervous system of a zebrafish larva, a preferred model organism for research. Motor neurons in the spinal cord can be seen in the researchers' images; at the same time, a single neuron with all its extensions is highlighted in another colour.


An observation by William Dempsey, post-doc in the group of ETH professor Periklis Pantazis, led to the new application. He worked with a special class of [fluorescent proteins](#) (see box) that change colour when irradiated with laser light of a specific wavelength. One such 'chameleon protein' is called Dendra 2, which normally emits green light when illuminated with blue light. The emission of Dendra 2 is however shifted into the red when it is irradiated by intensive violet or ultraviolet (UV) laser light.

## **Two-laser combination**

Dempsey and Pantazis specifically discovered that when Dendra 2 is irradiated by both a blue and a red laser at the same time, the protein's colour can also change to red. For this dual-colour illumination low intensity laser light is sufficient. In contrast to high intensity violet or ultraviolet (UV) irradiation it does not damage living cells.

ETH professor Pantazis and his colleagues then had an idea of how this finding could be deployed in light microscopy. Fluorescent proteins can

be used to make whole cells, precise cell structures or single molecules visible. For the first time, the ETH researcher's discovery permits a single cell or group of molecules located within a desirable part of a living organism to be highlighted with one colour, while all the other cells or molecules remain visible with another colour.



A color filter divides the laser light into two separate beams of different colors that overlap at a focal point (diagram). Credit: Patrick Helbling / ETH Zurich

## **Focus on one point**

The research group showed that when used individually, two different laser beams cannot change a chameleon protein's colour. But when the two beams are combined and directed in a way that they meet at a certain point on the object, then the proteins in focus change colour. In contrast, the proteins that are not activated at the same time by the two lasers retain their original colour.

The researchers have developed a simple and inexpensive colour filter, which can be used with the conventional confocal laser microscopes that are found in many biomedical research institutes. When mounted between the laser source and object, the filter divides the [laser light](#) into separate blue and red beams that are directed on to a small focal point on the object.

## **Examination of dynamic processes**

In the case of the zebrafish larva, which is transparent and therefore well suited for microscopy, the ETH researchers used Dendra 2 to colour neurons. They then focused the combined laser beam's focal point on the cell body of a single neuron in a live, anesthetized zebrafish. The local Dendra 2 molecules became red, spread throughout the entire cell and dyed the cell extensions. All other cells, even in the immediate vicinity of the targeted cell, remained green.

The ability to make individual neurons visible could be of great importance, for example, in the precise mapping of the brain, according to Pantazis. Since the method is suitable for individual cell targeting in living organisms, it could also be used to examine dynamic processes; for example, what happens to individual cells or a group of molecules when researchers treat an organism with active pharmaceutical ingredients. Embryo development could also be examined in more detail.

"Our method allows for a three-dimensional analysis in an elegant manner," explains Pantazis. "This is a very nice example of how you can take a result from basic research and use it to provide a solution for a technically challenging issue." Pantazis hopes the technique will be used more broadly in biomedical research in the future and is in talks with microscope manufacturers to implement this technology.

## Fluorescent proteins

Fluorescent proteins play a key role in [biomedical research](#). It is these proteins that researchers use to colour a cell structure or specific molecule, in order to make them visible under the microscope. In nature, these proteins are found in certain species of jellyfish and coral. Over time, researchers have created a rainbow colour palette of fluorescent proteins, which emit light within the visible spectrum. As a result, different cell structures can be made visible at the same time using fluorescent proteins with different colour emission. In addition, certain fluorescent proteins, such as the Dendra 2 protein that was used in this study, change their [colour](#) after irradiation with a certain wavelength of light.

**More information:** Dempsey WP, Georgieva L, Helbling PM, Sonay AY, Truong TV, Haffner M, Pantazis P: In vivo single cell labeling by confined primed conversion, *Nature Methods*, 18 May 2015, [DOI: 10.1038/nmeth.3405](https://doi.org/10.1038/nmeth.3405)

Provided by ETH Zurich

Citation: Chameleon proteins make individual cells visible (2015, May 19) retrieved 3 May 2024 from <https://phys.org/news/2015-05-chameleon-proteins-individual-cells-visible.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.