


# DNA double helix does double duty in assembling arrays of nanoparticles

May 25 2015



Scientists built octahedrons using ropelike structures made of bundles of DNA double-helix molecules to form the frames (a). Single strands of DNA attached at the vertices (numbered in red) can be used to attach nanoparticles coated with complementary strands. This approach can yield a variety of structures, including ones with the same type of particle at each vertex (b), arrangements with particles placed only on certain vertices (c), and structures with different

particles placed strategically on different vertices (d). Credit: Brookhaven National Laboratory

In a new twist on the use of DNA in nanoscale construction, scientists at the U.S. Department of Energy's (DOE) Brookhaven National Laboratory and collaborators put synthetic strands of the biological material to work in two ways: They used ropelike configurations of the DNA double helix to form a rigid geometrical framework, and added dangling pieces of single-stranded DNA to glue nanoparticles in place.

The method, described in the journal *Nature Nanotechnology*, produced predictable clusters and arrays of nanoparticles—an important step toward the design of materials with tailored structures and functions for applications in energy, optics, and medicine.

"These arrays of nanoparticles with predictable geometric configurations are somewhat analogous to molecules made of atoms," said Brookhaven physicist Oleg Gang, who led the project at the Lab's Center for Functional Nanomaterials (CFN), a DOE Office of Science User Facility. "While atoms form molecules based on the nature of their chemical bonds, there has been no easy way to impose such a specific spatial binding scheme on nanoparticles. This is exactly the problem that our method addresses."

Using the new method, the scientists say they can potentially orchestrate the arrangements of different types of nanoparticles to take advantage of collective or synergistic effects. Examples could include materials that regulate energy flow, rotate light, or deliver biomolecules.


"We may be able to design materials that mimic nature's machinery to harvest solar energy, or manipulate light for telecommunications

applications, or design novel catalysts for speeding up a variety of chemical reactions," Gang said.

The scientists demonstrated the technique to engineer nanoparticle architectures using an octahedral scaffold with particles positioned in precise locations on the scaffold according to the specificity of DNA coding. The designs included two different arrangements of the same set of particles, where each configuration had different optical characteristics. They also used the geometrical clusters as building blocks for larger arrays, including linear chains and two-dimensional planar sheets.

"Our work demonstrates the versatility of this approach and opens up numerous exciting opportunities for high-yield precision assembly of tailored 3D building blocks in which multiple nanoparticles of different structures and functions can be integrated," said CFN scientist Ye Tian, one of the lead authors on the paper.

## Details of assembly



A combination cryo-electron microscopy image of an octahedral frame with one gold nanoparticle bound to each of the six vertices, shown from three different angles. Credit: Brookhaven National Laboratory

This nanoscale construction approach takes advantage of two key characteristics of the DNA molecule: the twisted-ladder double helix shape, and the natural tendency of strands with complementary bases (the A, T, G, and C letters of the genetic code) to pair up in a precise way.

First, the scientists created bundles of six double-helix molecules, then put four of these bundles together to make a stable, somewhat rigid building material—similar to the way individual fibrous strands are woven together to make a very strong rope. The scientists then used these ropelike girders to form the frame of three-dimensional octahedrons, "stapling" the linear DNA chains together with hundreds of short complementary DNA strands.


"We refer to these as DNA origami octahedrons," Gang said.

To make it possible to "glue" nanoparticles to the 3D frames, the scientists engineered each of the original six-helix bundles to have one helix with an extra single-stranded piece of DNA sticking out from both ends. When assembled into the 3D octahedrons, each vertex of the frame had a few of these "sticky end" tethers available for binding with objects coated with complementary DNA strands.

"When nanoparticles coated with single strand tethers are mixed with the DNA origami octahedrons, the 'free' pieces of DNA find one another so the bases can pair up according to the rules of the DNA complementarity code. Thus the specifically DNA-encoded particles can find their correspondingly designed place on the octahedron vertices" Gang said.

The scientists can change what binds to each vertex by changing the DNA sequences encoded on the tethers. In one experiment, they

encoded the same sequence on all the octahedron's tethers, and attached strands with a complementary sequence to gold nanoparticles. The result: One gold nanoparticle attached to each of octahedron's six vertices.



By strategically placing tethers on particular vertices, the scientists used the octahedrons to link nanoparticles into one-dimensional chainlike arrays (left) and two-dimensional square sheets (right). Credit: Brookhaven National Laboratory

In additional experiments the scientists changed the sequence of some vertices and used complementary strands on different kinds of particles, illustrating that they could direct the assembly and arrangement of the particles in a very precise way. In one case they made two different arrangements of the same three pairs of particles of different sizes, producing products with different optical properties. They were even

able to use DNA tethers on selected vertices to link octahedrons end to end, forming chains, and in 2D arrays, forming sheets.

## Visualization of arrays

Confirming the particle arrangements and structures was a major challenge because the nanoparticles and the DNA molecules making up the frames have very different densities. Certain microscopy techniques would reveal only the particles, while others would distort the 3D structures.

To see both the particles and origami frames, the scientists used cryo-electron microscopy (cryo-EM), led by Brookhaven Lab and Stony Brook University biologist Huilin Li, an expert in this technique, and Tong Wang, the paper's other lead co-author, who works in Brookhaven's Biosciences department with Li. They had to subtract information from the images to "see" the different density components separately, then combine the information using single particle 3D reconstruction and tomography to produce the final images.

"Cryo-EM preserves samples in their near-native states and provides close to nanometer resolution," Wang said. "We show that cryo-EM can be successfully applied to probe the 3D structure of DNA-nanoparticle clusters."

These images confirm that this approach to direct the placement of [nanoparticles](#) on DNA-encoded vertices of molecular frames could be a successful strategy for fabricating novel nanomaterials.

**More information:** "Prescribed nanoparticle cluster architectures and low-dimensional arrays built using octahedral DNA origami frames": [dx.doi.org/10.1038/nnano.2015.105](https://doi.org/10.1038/nnano.2015.105)

Provided by Brookhaven National Laboratory

Citation: DNA double helix does double duty in assembling arrays of nanoparticles (2015, May 25) retrieved 2 May 2024 from <https://phys.org/news/2015-05-dna-helix-duty-arrays-nanoparticles.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--