

Research links prescribed burning to reduced tick populations

May 1 2015, by Molly Berg



Elizabeth Gleim, a former graduate student in the UGA Warnell School of Forestry and Natural Resources, checks tick sites.

In the spring, summer and early fall months, lone star ticks, the most common type of tick in Georgia, spike in activity and may transmit pathogens that could cause disease in humans. According to newly published University of Georgia research, tick populations can be managed through a technique called prescribed burning.

Elizabeth Gleim, a former [graduate student](#) in the Warnell School of Forestry and Natural Resources and co-author on the PLOS One article, performed the research over several years. During that time, she visited 21 plots in Georgia and Florida to collect tick samples monthly. Most of those plots had been subjected to long-term prescribed burning, a way to manage and control the growth of plant and animal species on a specific area of land.

Gleim, now a visiting assistant professor at Emory University's Oxford College, found that prescribed burning over a long period of time effectively reduced tick populations and changed the landscape of the burned areas.

"At the end of the day, prescribed fire on a long-term basis decimated tick populations," Gleim said. "It's exciting because by reducing tick populations, the technique indicated that it was reducing risk of tick-borne disease in humans while also enhancing ecosystem health."

Lone star [ticks](#) are known for transmitting a number of human pathogens that can cause disease. They include southern tick associated rash illness, known as STARI, and ehrlichiosis. Although ticks are present in Georgia and Florida year-round, they are a greater danger in the spring, summer and early fall when they are active.

Roy Berghaus, an epidemiologist in the UGA School of Veterinary Medicine, helped analyze the data on the ticks Gleim collected.



Lone star tick. Credit: Gerald Holmes, California Polytechnic State University at San Luis Obispo, Bugwood.org

"Any of the burned sites they looked at had a dramatic reduction in the number of ticks they were able to collect," Berghaus said. "Those tick sites that they couldn't do prescribed burning on had dramatic increases compared to the other burning sites."

At plots where there was no burning, tick counts were 10 times greater in the warm season than sites where prescribed burning occurred.

In total, over 47,000 ticks were collected among the 21 sites over two years. The most prominent of the ticks was the lone star tick, which is the most common tick in the Southeast.

Gleim stresses that repeat, not singular, prescribed burns will decrease tick populations.

"If you are in a fire-dependent or -tolerant ecosystem, then prescribed fire would be an effective way to deal with tick populations," she said. "It needs to be done on a long-term basis. Doing it once or twice would not permanently reduce tick populations. Prescribed fire needs to alter the vegetative structure of the forest in order to maintain the initial reduction of tick populations immediately following a fire. We were seeing that it was permanently decimating tick populations."



Elizabeth Gleim, a former graduate student in the Warnell School of Forestry

and Natural Resources, stands in front of a prescribed burning site used to reduce tick populations.

However, prescribed burning may not be an ideal solution if the land cannot support it.

"There are some limitations when it comes to prescribed burning," Berghaus said. "The land may be near a major highway, and it could cause an accident. In cases like that, there are some landscape and wildlife areas that need to take different measures."

Aside from [prescribed burning](#), there are ways individuals can keep safe from ticks year-round.

"The No. 1 thing is to perform prompt, careful full body checks after a visit into the forest," Gleim said. "While preventative measures such as wearing insect repellents with DEET or permethrin are wise, chances are some ticks will get past these defenses. The good news for most tick-borne pathogens is that it takes a tick several hours to transmit a pathogen. If you can remove the tick in that time, you'll be safe."

More information: "The Phenology of Ticks and the Effects of Long-Term Prescribed Burning on Tick Population Dynamics in Southwestern Georgia and Northwestern Florida." *PLoS ONE* 9(11): e112174. [DOI: 10.1371/journal.pone.0112174](https://doi.org/10.1371/journal.pone.0112174)

Provided by University of Georgia

Citation: Research links prescribed burning to reduced tick populations (2015, May 1) retrieved

26 April 2024 from <https://phys.org/news/2015-05-links-populations.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.