


Research team captures movement on Nepal earthquake fault rupture

May 6 2015, by Shayna Chabner Mckinney



The modeled slip on the fault is shown as viewed from above and indicated by the colors and contours within the rectangle. The peak slip in the fault exceeds 19.7 feet (6 meters). The ground motion measured with GPS is shown by the red and purple arrows and was used to develop the fault slip model. Aftershocks are indicated by red dots. Background color and shaded relief reflect regional variations in topography. The barbed lines show where the main fault reaches

Earth's surface. Credit: NASA/JPL-Caltech


Using a combination of satellite radar imaging data, GPS data measured in and near Nepal, and seismic observations from instruments around the world, Caltech and JPL scientists have constructed a preliminary picture of what happened below Earth's surface during the recent 7.8-magnitude Gorkha earthquake in Nepal.

The team's observations and models of the April 25, 2015 earthquake, produced through the [Advanced Rapid Imaging and Analysis \(ARIA\) project](#)—a collaboration between Caltech and JPL—include preliminary estimates of the slippage of the fault beneath Earth's surface that resulted in the deaths of thousands of people. In addition, the ARIA scientists have provided first responders and key officials in Nepal with information and maps that show block-by-block building devastation as well as measurements of ground movement at individual locations around the country.

"As the number of orbiting imaging radar and optical satellites that form the international constellation increases, the expected amount of time it takes to acquire an image of an impacted area will decrease, allowing for products such as those we have made for Nepal to become more commonly and rapidly available," says Mark Simons, professor of geophysics at Caltech and a member of the ARIA team. "I fully expect that within five years, this kind of information will be available within hours of a big disaster, ultimately resulting in an ability to save more lives after a disaster and to make assessment and response more efficient in both developed and developing nations."

Over the last five years, Simons and his colleagues in Caltech's Seismological Laboratory and at JPL have been developing the

approaches, infrastructure, and technology to rapidly and automatically use satellite-based observations to measure the movement of Earth's surface associated with earthquakes, volcanoes, landslides and other geophysical processes.



Color represents vertical movement and the scaled arrows indicate direction and magnitude of horizontal movement. Aftershocks are indicated by red dots. Background color and shaded relief reflect regional variations in topography. The barbed lines show where the main fault reaches Earth's surface. The main fault dives northward into the Earth below the Himalaya. Credit: NASA/JPL-Caltech

"ARIA is ultimately aimed at providing tools and data—for use by groups ranging from first responders, to government agencies, and individual scientists—that can help improve situational awareness, response, and recovery after many natural disasters," Simons says. "The same products also provide key observational constraints on our physical understanding of the underlying processes such as the basic physics controlling seismogenic behavior of major faults."

Provided by California Institute of Technology

Citation: Research team captures movement on Nepal earthquake fault rupture (2015, May 6)
retrieved 23 April 2024 from

<https://phys.org/news/2015-05-team-captures-movement-nepal-earthquake.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.