

# Analysis shows elements not acting in nature as previously modeled

June 16 2015



Workers on a cleanup site at DOE's Hanford Site in southeastern Washington State, one of several sites sampled for uranium and plutonium variations. Credit: U.S. Department of Energy

Knowing how a chemical in soil reacts and transforms over time in

response to neighboring elements, weather and heat is essential in determining whether that chemical is hazardous. This is especially important when that chemical is radioactive.

In a collaborative, international effort led by Los Alamos National Laboratory, researchers determined the speciation of uranium and [plutonium](#) pulled from soil, concrete and water reservoir sediment from six locations in the U.S., Ukraine and Russia. Speciation is the chemical form or compound in which an element can appear, for example as nitrogen can form species of nitrate, nitrite or ammonia, among others.

"These results demonstrate the complexities of actinides in the environment and suggest that our models will not always accurately predict their long term fate and transport," said Robert Roback, a Los Alamos Earth and Environmental Sciences Division deputy group leader and coauthor on the paper.

## **Novel results, environmental implications**

Results clearly identified novel aspects of uranium and plutonium environmental chemistry beyond those previously reported and expected based on thermodynamics. In some samples, actinide-bearing materials that were within millimeters of each other for decades still exhibit significantly different speciation.

These results, the authors note, demonstrate complexities of actinides in the environment that are likely linked to the initial depositional form as well as to their subsequent mobilization and deposition. "These results do not necessarily imply increased environmental risk, however, because some forms are quite stable and have low solubility," Roback said. The results imply that to understand predict uranium and plutonium behavior in the environment, the analysis must consider site conditions, he said.

The researchers published their findings in a cover article, "Multiscale Speciation of U and Pu at Chernobyl, Hanford, Los Alamos, McGuire AFB, Mayak, and Rocky Flats" recently published by the American Chemical Society's journal *Environmental Science and Technology*.

The researchers identified four types of both equilibrium species (mostly stabilized) and unpredicted novel types of these actinides unassociated with the source, which helped elucidate if and how the various forms of the elements may have been transformed by chemical reactions in the environment.

## **Hazardous materials transformation**

The chemical speciation of potentially hazardous subsurface contaminants is crucial to assessing environmental risk and remediation, because it determines the contaminants' transport and toxicology. Understanding the behavior of uranium and plutonium contamination is imperative since not knowing the behavior is a significant impediment to environmental restoration at legacy nuclear weapons' production and testing sites plus reactor accident locations.

Information obtained in the study will help researchers validate models that calculate safe residual levels, reducing the amount of hazardous material requiring removal and disposition.

The scientists retrieved samples from Chernobyl (nuclear power plant) and Hanford Site (plutonium reactors), Los Alamos (Manhattan Project-era waste), McGuire Air Force Base (nuclear warhead fire), Mayak (irradiated fuel processing waste) and Rocky Flats Environmental Technology Site (weapons' fissile material fabrication).

Materials from some of these sites involved high temperatures, melting, fires and explosions. In some cases, uranium and plutonium were known

or suspected to have been released in acidic solutions that reacted with soil or concrete components. The goal was to determine the range of behavior in contamination and releases caused by [chemical reactions](#).

Using synchrotron microprobes, researchers scanned soil to locate actinide hotspots and associated elements before performing more detailed analysis. Measurements were performed on beamlines at the Stanford Synchrotron Radiation Lightsource. Quantitative electron probe microanalysis of Chernobyl samples was carried out at the V.G. Khlopin Radium Institute.

Higher temperatures (such as in the Chernobyl explosion and fire) promoted reactions and materials mixing. Samples revealed that in some cases particles melded without significant mixing, whereas other actinide particles intermingled with other metals, air and water to give a more homogenous mixture.

In a soil sample collected from Hanford, two distinct plutonium populations occur. One population, consisting of 20-micron cubes with uniformly low plutonium concentrations, was thought to have formed during initial deposition of the plutonium carried in tributyl phosphate. Interestingly, a second population contains large plutonium concentrations as surface precipitates. The researchers suggest that this second population is evidence for plutonium mobilization and deposition onto the previously existing cubes found at Hanford.

**More information:** "Multiscale Speciation of U and Pu at Chernobyl, Hanford, Los Alamos, McGuire AFB, Mayak, and Rocky Flats." *Environ. Sci. Technol.*, 2015, 49 (11), pp 6474–6484 [DOI: 10.1021/es506145b](#)

Provided by Los Alamos National Laboratory

Citation: Analysis shows elements not acting in nature as previously modeled (2015, June 16)  
retrieved 2 May 2024 from

<https://phys.org/news/2015-06-analysis-elements-nature-previously.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--