

Researchers discover why 'blue hook' stars are unusually hot

June 23 2015



The crowded centre of the Omega Centauri globular cluster. Credit: Hubble Space telescope, NASA

Astronomers have solved a mystery over small, unusually hot blue stars, 10 times hotter than our Sun, that are found in the middle of dense star clusters.

The international team found the so-called blue hook [stars](#) throw off

their cool outer layers late in life because they are rotating so rapidly, making them more luminous than usual.

"We've solved an old puzzle. These stars are only half the mass of our Sun yet we could not explain how they became so luminous," said team member Dr Antonino Milone, from ANU Research School of Astronomy and Astrophysics.


"As the star was forming billions of years ago from a disc of gas in the congested centre of the [star cluster](#), another star or stars must have collided with the disc and destroyed it."

The research, published in *Nature*, gives new insights into star formation in the early Universe in the crowded centres of clusters. Star clusters are rare environments in the Universe, in which many stars are born at the same time.

The team studied the globular cluster Omega Centauri, the only cluster visible to the naked eye, which contains around 10 million stars in close proximity to one another.

The model shows the formation of stars in clusters do not all form at once, said co-author Dr Aaron Dotter, also from ANU Research School of Astronomy and Astrophysics.

"These blue stars must form in a second generation of [star formation](#)," he said. "Our new explanation is quite simple, and it hangs together really nicely."



Omega Centauri globular cluster. Credit:ESO/INAF-VST/OmegaCAM

Usually the large disc of ionised gas around a newly-forming star locks its rotation through magnetic effects. For the progenitors of blue hook stars, however, an early destruction of its disc allows the stars to spin up as the gas comes together to form a star.

Because its high rotation rate partially balances the inward force of gravity, the star consumes its hydrogen fuel more slowly and evolves differently throughout its life.

The blue hook phase of its life occurs after more than 10 billion years, when the star has consumed nearly all its hydrogen and begins burning the hotter fuel helium. The different evolution processes leave it with a heavier core which burns brighter than typical helium-burning stars.



Artist's impression of a protostellar disc disrupted by a collision. Credit: Marco Galliani, INAF

More information: "Rapidly rotating second-generation progenitors for the 'blue hook' stars of ω Centauri" *Nature* (2015) [DOI: 10.1038/nature14516](https://doi.org/10.1038/nature14516)

Provided by Australian National University

Citation: Researchers discover why 'blue hook' stars are unusually hot (2015, June 23) retrieved 1 May 2024 from <https://phys.org/news/2015-06-blue-stars-unusually-hot.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.