


# Giant galaxy is still growing

June 25 2015

---



The huge halo around giant elliptical galaxy Messier 87 appears on this very deep

image. An excess of light in the top-right part of this halo, and the motion of planetary nebulae in the galaxy, are the last remaining signs of a medium-sized galaxy that recently collided with Messier 87. The image also reveals many other galaxies forming the Virgo Cluster, of which Messier 87 is the largest member. In particular, the two galaxies at the top right of the frame are nicknamed 'the Eyes'. Credit: Chris Mihos (Case Western Reserve University)/ESO

New observations with ESO's Very Large Telescope have revealed that the giant elliptical galaxy Messier 87 has swallowed an entire medium-sized galaxy over the last billion years. For the first time a team of astronomers has been able to track the motions of 300 glowing planetary nebulae to find clear evidence of this event and also found evidence of excess light coming from the remains of the totally disrupted victim.

Astronomers expect that [galaxies](#) grow by swallowing smaller galaxies. But the evidence is usually not easy to see—just as the remains of the water thrown from a glass into a pond will quickly merge with the pond water, the stars in the infalling galaxy merge in with the very similar stars of the bigger galaxy leaving no trace.

But now a team of astronomers led by PhD student Alessia Longobardi at the Max-Planck-Institut für extraterrestrische Physik, Garching, Germany has applied a clever observational trick to clearly show that the nearby [giant elliptical galaxy](#) Messier 87 merged with a smaller spiral galaxy in the last billion years.

"This result shows directly that large, luminous structures in the Universe are still growing in a substantial way—galaxies are not finished yet!" says Alessia Longobardi. "A large sector of Messier 87's outer halo now appears twice as bright as it would if the collision had not taken place."

Messier 87 lies at the centre of the Virgo Cluster of galaxies. It is a vast ball of stars with a total mass more than a million million times that of the Sun, lying about 50 million light-years away.

Rather than try to look at all the stars in Messier 87—there are literally billions and they are too faint and numerous to be studied individually—the team looked at planetary nebulae, the glowing shells around ageing stars. Because these objects shine very brightly in a specific hue of aquamarine green, they can be distinguished from the surrounding stars. Careful observation of the light from the nebulae using a powerful spectrograph can also reveal their motions.

Just as the water from a glass is not visible once thrown into the pond—but may have caused ripples and other disturbances that can be seen if there are particles of mud in the water—the motions of the planetary nebulae, measured using the FLAMES spectrograph on the Very Large Telescope, provide clues to the past merger.

"We are witnessing a single recent accretion event where a medium-sized galaxy fell through the centre of Messier 87, and as a consequence of the enormous gravitational tidal forces, its stars are now scattered over a region that is 100 times larger than the original galaxy!" adds Ortwin Gerhard, head of the dynamics group at the Max-Planck-Institut für extraterrestrische Physik, Garching, Germany, and a co-author of the new study.

The team also looked very carefully at the light distribution in the outer parts of Messier 87 and found evidence of extra light coming from the stars in the galaxy that had been pulled in and disrupted. These observations have also shown that the disrupted galaxy has added younger, bluer stars to Messier 87, and so it was probably a star-forming [spiral galaxy](#) before its merger.

"It is very exciting to be able to identify stars that have been scattered around hundreds of thousands of light-years in the halo of this galaxy—but still to be able to see from their velocities that they belong to a common structure. The green planetary nebulae are the needles in a haystack of golden stars. But these rare needles hold the clues to what happened to the [stars](#)," concludes co-author Magda Arnaboldi (ESO, Garching, Germany).

**More information:** This research was presented in a paper entitled "The build-up of the cD halo of M87—evidence for accretion in the last Gyr", by A. Longobardi et al., to appear in the journal *Astronomy & Astrophysics Letters* on 25 June 2015.

[dx.doi.org/10.1051/0004-6361/201526282](https://doi.org/10.1051/0004-6361/201526282) , [arxiv.org/abs/1504.04369](https://arxiv.org/abs/1504.04369)

Provided by ESO

Citation: Giant galaxy is still growing (2015, June 25) retrieved 25 April 2024 from <https://phys.org/news/2015-06-giant-galaxy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.