

High vitamin C levels are required to overcome photo-inhibition in plants

June 24 2015




Figure 1. AtPHT4;4 is a vitamin C transporter in Arabidopsis. (Left) Phylogenetic tree of the plant PHT4 family. (Right) Vitamin C uptake by proteoliposomes containing AtPHT4;4 or no AtPHT4;4.

Vitamin C (ascorbate) is an antioxidant and coenzyme for a number of metabolic reactions in living organisms. In plant chloroplasts, high vitamin C levels are required to overcome photo-inhibition caused by

strong light.

Although [vitamin](#) C is synthesized in the mitochondria, the molecular mechanisms underlying vitamin C [transport](#) into [chloroplasts](#) are poorly understood.

Now, Takaaki Miyaji, Yoshinori Moriyama, and Jian Feng Ma at Okayama University, together with Takashi Kuromori at RIKEN and colleagues, have shown that AtPHT4;4, a member of the phosphate transporter 4 family of *Arabidopsis thaliana*, functions as a vitamin C transporter.

The team found that proteoliposomes containing purified AtPHT4;4 protein exhibited membrane potential ($\Delta\Psi$)- and Cl⁻—dependent vitamin C uptake (Fig. 1). AtPHT4;4 protein was expressed abundantly in the chloroplast envelope membranes (Fig. 2). Knocking out AtPHT4;4 resulted in decreased levels of vitamin C in the chloroplasts. The heat dissipation process of excessive energy during photosynthesis also decreased in the mutants.

These results indicate that AtPHT4;4 protein is a vitamin C transporter at the envelope membranes of chloroplasts, which is required for tolerance to strong light stress (Fig. 3). This is the first report regarding the identification of a vitamin C transporter in plants.

This research could help guide the development of heritable transporter genetic modification technology, which may provide a means of developing photo-inhibition tolerant plants.




Figure 2. AtPHT4;4 is abundantly expressed in the envelope membranes of chloroplasts from the leaves of Arabidopsis. The fluorescence signals of AtPHT4;4 and chlorophyll are shown in green and magenta, respectively.




Figure 3. The mechanism of vitamin C transport in chloroplast. Upon strong light, the PHT4;4 at the envelope membranes takes up vitamin C from mitochondria, which is required for photostress tolerance.

More information: "AtPHT4;4 is a chloroplast-localized ascorbate transporter in Arabidopsis." *Nature Communications* 6, Article number: 5928 DOI: [10.1038/ncomms6928](https://doi.org/10.1038/ncomms6928)

Provided by Okayama University

Citation: High vitamin C levels are required to overcome photo-inhibition in plants (2015, June 24) retrieved 1 June 2024 from <https://phys.org/news/2015-06-high-vitamin-required-photo-inhibition.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.