


# Hubble detects 'sunscreen' layer on distant planet

June 11 2015, by Karen Northon



WASP-33b's stratosphere was detected by measuring the drop in light as the planet passed behind its star (top). Temperatures in the low stratosphere rise because of molecules absorbing radiation from the star (right). Without a stratosphere, temperatures would cool down at higher altitudes (left). Credit: NASA/Goddard

NASA's Hubble Space Telescope has detected a stratosphere, one of the primary layers of Earth's atmosphere, on a massive and blazing-hot exoplanet known as WASP-33b.

The presence of a [stratosphere](#) can provide clues about the composition of a planet and how it formed. This atmospheric layer includes molecules that absorb ultraviolet and visible light, acting as a kind of 'sunscreen' for the planet it surrounds. Until now, scientists were uncertain whether these molecules would be found in the atmospheres of large, extremely hot planets in other star systems. These findings will appear in the June 12 issue of the *Astrophysical Journal*.

"Some of these planets are so hot in their upper atmospheres, they're essentially boiling off into space," said Avi Mandell, a planetary scientist at NASA's Goddard Space Flight Center in Greenbelt, Md., and a co-author of the study. "At these temperatures, we don't necessarily expect to find an atmosphere that has molecules that can lead to these multilayered structures."

In Earth's atmosphere, the stratosphere sits above the troposphere—the turbulent, active-weather region that reaches from the ground to the altitude where nearly all clouds top out. In the troposphere, the temperature is warmer at the bottom—ground level—and cools down at higher altitudes.

The stratosphere is just the opposite. In this layer, the temperature increases with altitude, a phenomenon called temperature inversion. On Earth, temperature inversion occurs because ozone in the stratosphere absorbs much of the sun's [ultraviolet radiation](#), preventing it from reaching the surface, protecting the biosphere, and therefore warming the stratosphere instead.

Similar temperature inversions occur in the stratospheres of other planets

in our solar system, such as Jupiter and Saturn. In these cases, the culprit is a different group of molecules called hydrocarbons. Neither ozone nor hydrocarbons, however, could survive at the high temperatures of most known exoplanets, which are planets outside our solar system. This leads to a debate as to whether stratospheres would exist on them at all.

Using Hubble, the researchers have settled this debate by identifying a temperature inversion in the atmosphere of WASP-33b, which has about four-and-a-half times the mass of Jupiter. Team members also think they know which molecule in WASP-33b's atmosphere caused the inversion—titanium oxide.

"These two lines of evidence together make a very convincing case that we have detected a stratosphere on an exoplanet," said Korey Haynes, lead author of the study. Haynes was a graduate student at George Mason University in Fairfax, Va., and was working at Goddard with Mandell when the research was conducted.

The researchers analyzed observations made with Hubble's Wide Field Camera 3 by co-author Drake Deming at the University of Maryland in College Park. Wide Field Camera 3 can capture a spectrum of the near-infrared region where the signature for water appears. Scientists can use the spectrum to identify water and other gases in a distant planet's atmosphere and determine its temperature.

Haynes and her colleagues used the Hubble observations, and data from previous studies, to measure emission from water and compare it to emission from gas deeper in the atmosphere. The team determined that emission from water was produced in the stratosphere at about 6,000 degrees Fahrenheit. The rest of the emission came from gas lower in the atmosphere that was at a temperature about 3,000 degrees Fahrenheit.

The team also presented the first observational evidence that

WASP-33b's atmosphere contains titanium oxide, one of only a few compounds that is a strong absorber of visible and ultraviolet radiation and capable of remaining in gaseous form in an [atmosphere](#) as hot as this one.

"Understanding the links between stratospheres and chemical compositions is critical to studying atmospheric processes in exoplanets," said co-author Nikku Madhusudhan of the University of Cambridge, United Kingdom. "Our finding marks a key breakthrough in this direction."

**More information:** "Spectroscopic Evidence for a Temperature Inversion in the Dayside Atmosphere of the Hot Jupiter WASP-33b," Korey Haynes et al., 2015, *Astrophysical Journal*, Preprint: [arxiv.org/abs/1505.01490](https://arxiv.org/abs/1505.01490)

Provided by NASA's Goddard Space Flight Center

Citation: Hubble detects 'sunscreen' layer on distant planet (2015, June 11) retrieved 1 May 2024 from <https://phys.org/news/2015-06-hubble-sunscreen-layer-distant-planet.html>

|  |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|