

# Scientists help public avoid health risks of toxic blue-green algae

June 15 2015

---



*Dolichospermum crissum*, one of the most problematic cyanobacteria in Alberta lakes. Credit: FlowCAM photo, courtesy Mark Graham

As we approach the hottest days of the summer season, toxic algae are starting to bloom in lakes around the world. Increases in atmospheric temperature combined with land nutrients are promoting the growth of harmful cyanobacteria—blue-green algae—in these bodies of water, damaging for not only the associated ecosystems but also potentially for the people who recreationally frequent the lakes.

University of Alberta scientists work to mitigate the risks to human

health with the use of advanced flow cytometry, a technique they've developed and have been using for the past three years.

"Our ability to rapidly detect the onset of outbreaks of potentially [toxic cyanobacteria](#) provides timely information to the public regarding the major lakes throughout the province," says Professor Rolf Vinebrooke (Biological Sciences). "If the cell counts exceed the World Health Organization guideline of 100,000 cells per millilitre, we notify Alberta Health officials, who then post advisories at the lake in question."

Vinebrooke and his colleagues use a "FlowCAM" unit, manufactured by Fluid Imaging Technologies, for early detection of [cyanobacteria](#) in lakes. Their protocol is the first adaptation in North America of the protocol used by oceanographers to warn of potential shellfish poisoning events produced by toxic marine algae. A sample of lake water is simply allowed to flow through the cytometer, which is equipped with imaging software that detects and measures more than 25 different optical properties of each cell. At the same time, the software constructs a collage of all the images and then matches each with a known cyanobacterium using a searchable electronic library of more than 100 species, developed by research associate Dr. Mark Graham.

"The next step is to detect trends in our cyanobacterial-monitoring data as related to measured environmental conditions," continues Vinebrooke. "This will enable us to generate both predictive models for blue-green algal blooms and testable hypotheses regarding the as-yet unknown set of factors that cause these outbreaks in the first place."



Cyanobacterium blooming in June 2015, Gainford Beach on Isle Lake, west of Edmonton, Alberta. Credit: Photo courtesy Leyna Tsui

Vinebrooke recently collaborated with colleagues from McGill University to quantitatively synthesize long-term worldwide trends of cyanobacterial growth. The scientists found that, as humans have altered the global climate over the past two centuries, there has been a disproportionate growth of blue-green algae relative to other phytoplankton, with particularly accelerated growth in the past 70 years.

"Excess inputs of nutrients into lakes combined with higher water temperatures due to climate warming appear to be two of the major

causes for the outbreaks," he says. "We recommend reducing the use of fertilizers and fossil fuels to help slow the global acceleration of this cyanobacterial dominance."

These findings were recently published in the scientific journal *Ecology Letters*.

Provided by University of Alberta

Citation: Scientists help public avoid health risks of toxic blue-green algae (2015, June 15)  
retrieved 2 May 2024 from

<https://phys.org/news/2015-06-scientists-health-toxic-blue-green-algae.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--