

The Southeast Pacific produces more nitrous oxide than previously thought

June 22 2015



View over the tropical Southeast Pacific from the research vessel METEOR. As new data show the region is a strong source of natural nitrous oxide. Credit: Kerstin Nachtigall, GEOMAR


In addition to carbon dioxide there are plenty of other greenhouse gases.

Nitrous oxide is one of them. However, a global assessment of emissions from the oceans is difficult because the measurement methods used so far have only allowed rough estimates. Using a new technology for continuous measurements, researchers of the GEOMAR and the Kiel University have now discovered that nitrous oxide emissions from the Southeast Pacific are much higher than previously thought. They publish their data in the international journal *Nature Geoscience*.

Originally it became famous as an anesthetic gas used by dentists. However, laughing gas, or chemically correct nitrous oxide, is also found in large quantities in nature and has serious effects on climate: In the lower atmosphere it is a strong [greenhouse gas](#), and in higher layers of the atmosphere it contributes indirectly to the destruction of ozone. "A global assessment of marine [nitrous oxide emissions](#) is, however, difficult because we do not know exactly where and how much nitrous oxide is produced," says marine chemist Damian L. Arévalo-Martínez from GEOMAR Helmholtz Centre for Ocean Research Kiel. Together with colleagues from GEOMAR and the University of Kiel (CAU), he presents new data in the international scientific journal *Nature Geoscience*, showing that the Southeast Pacific has been significantly underestimated as a source of nitrous oxide.

The published data are based on three expeditions of the German research vessel METEOR, which took place off Peru between November 2012 and March 2013. Together, the Kiel-based collaborative research centre "SFB 754" and the SOPRAN project have studied the extensive oxygen minimum zone (OMZ) off Peru since 2008. "In that area, like on the eastern boundaries of other tropical oceans, nutrient-rich waters from deeper water layers are transported to the surface," explains co-author Prof. Dr. Hermann Bange, also from GEOMAR. This results in intense plankton growth close to the surface, which upon death, sinks on the water column.

When microorganisms decompose this biomass, they thereby consume more oxygen than can be supplied by surrounding waters and thus the oxygen concentration decreases. Of all the tropical OMZs the one in the Pacific is the largest. "We know that oxygen depletion also affects the nitrogen cycle and favors the production of nitrous oxide," says Damian L. Arévalo-Martínez. However, previous measurements allowed only rough estimations of its release to the atmosphere.



The graph shows in the dark blue to violet tones the OMZs in the tropical oceans. They play a major role in the nitrogen cycle. For the SFB 754 they are the main working areas. Credit: SFB754

During the expeditions in 2012 and 2013, the researchers were able to measure nitrous oxide concentrations continuously from the METEOR for the first time. "Before, the ship had to stop every few miles. We took seawater samples, analyzed them and had data for one point only," explains Professor Bange. He adds: "With the new continuous measurement method we get a lot more data, which also enable us to make better extrapolations for the whole area."

These new data include, among others, the highest ever measured nitrous oxide concentrations in marine surface waters. "Our estimates revealed that each year about 0.3 to 1.4 megatons of nitrous oxide are emitted to the atmosphere from the oxygen minimum zone in the tropical Southeast Pacific. This is up to a fifth of the estimated marine [nitrous oxide](#) emissions worldwide and significantly exceeds similar upwelling areas in other tropical oceans," says Arévalo-Martínez.

The investigation of [greenhouse gas emissions](#) is also important to anticipate future changes in time and to assess their impact on climate. "The OMZs have expanded in recent years, but before we can see an increase or decrease of greenhouse gas emissions, we need to know the actual state," says Professor Bange.

More information: Arévalo-Martínez, D. L., A. Kock, C. R. Löscher, R. A. Schmitz, H. W. Bange (2015): Massive nitrous oxide emissions from the tropical South Pacific Ocean. *Nature Geoscience*, [dx.doi.org/10.1038/NGEO2469](https://doi.org/10.1038/NGEO2469)

Provided by Helmholtz Association of German Research Centres

Citation: The Southeast Pacific produces more nitrous oxide than previously thought (2015, June 22) retrieved 26 April 2024 from <https://phys.org/news/2015-06-southeast-pacific-nitrous-oxide->

[previously.html](#)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.