

Astronomers bring a new hope to find 'Tatooine' planets

July 16 2015, by Blaine Friedlander



Sibling suns – made famous in the "Star Wars" scene where Luke Skywalker gazes toward a double sunset – and the planets around them may be more common than we've thought, and Cornell astronomers are presenting new ideas on how to find them.

Astronomers could discover a plethora of <u>planets</u> around <u>binary star</u> <u>systems</u> (stars that rotate around each other) by measuring with high precision how stars move around each other, looking for disturbances exerted by possible exoplanets. So explains new research, "Survival of



Planets Around Shrinking Stellar Binaries," published in the *Proceedings of the National Academy Sciences*, July 9, by Diego J. Munoz, Cornell postdoctoral researcher, and Dong Lai, professor of astronomy, in the College of Arts and Sciences.

What once was fictional as young Skywalker saw the double suns from Tatooine is astronomical reality four decades later. Normal binary suns orbit each other every eight to 100 days, and the Kepler telescope easily can detect those exoplanets (planets outside of our own solar system) as the planets transit (move across) each <u>sun</u>.

Trouble starts in compact binary sun systems – where sibling suns move closer together – making it difficult for the most advanced telescopes to find them. Essentially, for Kepler and other telescopes, the planetary orbital plane of these double suns and their accompanying planets might be out of whack – or misaligned – rendering them invisible to us. "The current observational strategy inevitably misses a population of Tatooine planets, but future observations may reveal their existence," said Munoz.

NASA's Kepler telescope is a heliocentric (it orbits the sun) spacecraft that monitors star brightness in a Milky Way region near the constellation Cygnus, the swan. Measuring photons, Kepler detects lower light values – and thus, a planetary transit.

Munoz explains that suns in the close binary system likely were once standard systems that have lost energy and shrunk, bringing the suns closer together. As the sibling sun's distance decreases, the orbits of that system's planets become misaligned, rendering it impossible for the Kepler telescope to detect planets – which no longer cross in the front of the suns.

Munoz and Lai suggest scouting for exoplanet-caused disturbances for compact binary star systems, to determine a new population of



circumbinary planets. Said Munoz: "Since this type of 'compact' binary is very common, it had been very puzzling that no planets had been detected."

More information: Survival of Planets Around Shrinking Stellar Binaries, *PNAS*, 10.1073/pnas.1505671112 . On *Arxiv*: arxiv.org/abs/1505.05514

Provided by Cornell University

Citation: Astronomers bring a new hope to find 'Tatooine' planets (2015, July 16) retrieved 1 May 2024 from <u>https://phys.org/news/2015-07-astronomers-tatooine-planets.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.