

Researchers discover how bacteria sweet-talk their way into plants

July 9 2015



Root nodules, each containing billions of Rhizobiaceae bacteria. Credit: Public Domain

A University of Otago researcher is part of an international team of scientists that has discovered how legumes, which include important

agricultural species such as white clover, are able to tell helpful and harmful invading bacteria apart.

Their study, which changes the understanding of carbohydrates as signal molecules, is newly published in the leading international journal *Nature*. The research has implications for improving the understanding of how other plant and animal species interact with [bacteria](#) in their environment and defend themselves against hostile infections.

Clover and other legumes form a unique symbiotic relationship with bacteria known as rhizobia, which they allow to infect their roots. This leads to root nodules being formed in which the bacteria convert nitrogen from the air into ammonia that the plant can use for growth.

Exactly how these plants are able to distinguish and welcome in compatible [rhizobia](#) for this self-fertilising activity—while halting infection by incompatible bacteria—has been a mystery.

Now Otago's Professor Clive Ronson and colleagues in a research team led by Professor Jens Stougaard in Denmark at the Centre for Carbohydrate Recognition and Signalling (CARB) have determined how legumes perceive and distinguish compatible bacteria based on the exopolysaccharides featuring on the invading cells' surfaces.

Professor Ronson says that exopolysaccharides, along with other carbohydrates, are found on the cell surfaces of all organisms.

"Bacteria display a wide selection of polysaccharides, and many of these are important for cell-to-cell interactions, immune evasion, pathogenesis, biofilm formation and colonisation of ecological niches," he says.

Using an interdisciplinary approach involving plant and microbial genetics, biochemistry and [carbohydrate](#) chemistry, the CARB

researchers have identified the first known exopolysaccharide receptor gene, called Epr3.

They found that a membrane-bound receptor kinase (a type of enzyme) encoded by the Epr3 gene binds directly with exopolysaccharides and regulates [beneficial bacteria](#)'s passage through the plant's epidermal cell layer.

"Now that we have identified this key mechanism that allows a host organism to distinguish friendly bacteria from those that cause disease, this opens up exciting new research avenues across a number of fields," says Professor Ronson.

Microbiome studies in plants, animals and humans are some of the areas that will benefit from the new discovery. The mechanism governing microbiota colonisation of hosts is poorly understood and the identification of an exopolysaccharide receptor is likely to inspire new approaches to understand the interaction between multicellular organisms and microbes, he says.

More information: "Receptor-mediated exopolysaccharide perception controls bacterial infection." *Nature* (2015) [DOI: 10.1038/nature14611](https://doi.org/10.1038/nature14611)

Provided by University of Otago

Citation: Researchers discover how bacteria sweet-talk their way into plants (2015, July 9) retrieved 6 May 2024 from <https://phys.org/news/2015-07-bacteria-sweet-talk.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
