


# A dark matter bridge in our cosmic neighborhood

July 14 2015




The figure shows the current stream of galaxies - the flow along in the cosmic super-highway and on the bridge to Virgo, in the region around the Milky Way, Andromeda and Centaurus A

By using the best available data to monitor galactic traffic in our neighborhood, Noam Libeskind from the Leibniz Institute for Astrophysics Potsdam (AIP) and his collaborators have built a detailed

map of how nearby galaxies move. In it they have discovered a bridge of dark matter stretching from our Local Group all the way to the Virgo cluster—a huge mass of some 2,000 galaxies roughly 50 million light-years away, that is bound on either side by vast bubbles completely devoid of galaxies. This bridge and these voids help us understand a 40 year old problem regarding the curious distribution of dwarf galaxies.

These [dwarf galaxies](#) are often found swarming around larger hosts like our own Milky Way. Since they are dim they are hard to detect, and are thus found almost exclusively in our cosmic neighborhood. A particularly fascinating aspect of their existence is that near the Milky Way and at least two of our closest neighbors—the Andromeda and Centaurus A galaxies—these satellites don't just fly around randomly, but are instead compressed on to vast, flat, possibly spinning, planes. Such structures are not a naive outcome of the cold [dark matter](#) model that most cosmologists believe is responsible for how the universe forms galaxies. These structures are thus a challenge to the current doctrine.

One possibility is that these [small galaxies](#) echo the geometry of structure on much greater scales. "This is the first time we have had observational verification that large filamentary super highways are channeling dwarf galaxies across the cosmos along magnificent bridges of dark matter," Libeskind says. This cosmic "super highway" gives the speeding satellites an off ramp along which they can be beamed towards the Milky Way, Andromeda and Centaurus A. "The fact that this galactic bridge can affect the dwarf galaxies around us is impressive, given the difference in scale between the two: the planes of dwarfs are around 1 percent of the size of the galactic bridge to Virgo."



Current stream of galaxies (detailed).

**More information:** "Planes of Satellite Galaxies and the Cosmic Web," Noam Libeskind et al., 2015, *Monthly Notices of the Royal Astronomical Society* [mnras.oxfordjournals.org/content/452/1/1052](https://mnras.oxfordjournals.org/content/452/1/1052) ,  
*Arxiv:* [arxiv.org/abs/1503.05915](https://arxiv.org/abs/1503.05915)

Provided by Leibniz Institute for Astrophysics Potsdam

Citation: A dark matter bridge in our cosmic neighborhood (2015, July 14) retrieved 28 April

2024 from <https://phys.org/news/2015-07-dark-bridge-cosmic-neighborhood.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.