


Past and present sea levels in the Chesapeake Bay Region, USA

July 28 2015



This is a time series of the Blackwater River valley. Top: Intact marsh surveyed from AD 1902 to AD 1904 and presented in a 7.5" USGS topographic map from AD 1905; dark blue hatching around the Blackwater valley is tidal marsh; light blue pattern is freshwater swamp. Middle: Initiation of major ponding seen in an aerial photograph from AD 1938. Bottom: Coalesced ponds forming the informal 'Lake Blackwater' in satellite imagery from AD 2007. Credit: *GSA Today*, DeJong et al.,

In a new article for *GSA Today*, authors Benjamin DeJong and colleagues write that sea-level rise (3.4 mm/yr) is faster in the Chesapeake Bay region than any other location on the Atlantic coast of North America, and twice the global average (1.7 mm/yr). They have found that dated interglacial deposits suggest that relative sea levels in the Chesapeake Bay region deviate from global trends over a range of timescales.

According to DeJong and [colleagues](#), "Glacio-isostatic adjustment of the land surface from loading and unloading of continental ice is likely responsible for these deviations, but our understanding of the scale and timeframe over which isostatic response operates in this region remains incomplete because dated [sea-level](#) proxies are mostly limited to the Holocene and to deposits 80 ka or older."

To better understand glacio-isostatic control over past and present relative sea level, DeJong and colleagues applied a suite of dating methods to the stratigraphy of the Blackwater National Wildlife Refuge, one of the most rapidly subsiding and lowest-elevation surfaces bordering Chesapeake Bay. Their data indicate that the region was submerged at least for portions of marine isotope stage (MIS) 3 (about 30 to 60 million years ago), although, they note, multiple proxies suggest that [global sea level](#) was 40 to 80 meters lower than today.

Today, MIS 3 deposits are above sea level because they were raised by the Last Glacial Maximum forebulge, but decay of that same forebulge is causing ongoing subsidence. "These results," they write, "suggest that glacio-isostasy controlled relative sea level in the mid-Atlantic region for tens of thousands of years following retreat of the Laurentide Ice Sheet and continues to influence relative sea level in the region."

The study finds that isostatically driven subsidence of the Chesapeake Bay region will continue for millennia, exacerbating the effects of global

[sea-level rise](#) and impacting the region's large population centers and valuable coastal natural resources.

More information: Pleistocene relative sea levels in the Chesapeake Bay region and their implications for the next century, Benjamin D. DeJong, Paul R. Bierman, Wayne L. Newell, Tammy M. Rittenour, Shannon A. Mahan, Greg Balco, and Dylan H. Rood. Pages 4–10; [DOI: 10.1130/GSATG223A.1](#)

Provided by Geological Society of America

Citation: Past and present sea levels in the Chesapeake Bay Region, USA (2015, July 28)
retrieved 30 April 2024 from <https://phys.org/news/2015-07-sea-chesapeake-bay-region-usa.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--