

Spintronics just got faster

July 20 2015



This the 2-dimensional Ultrafast UV spectroscopy set-up at EPFL's Laboratory of Ultrafast Spectroscopy, used to carry out the measurements in this study. Credit: Alain Herzog/EPFL

In a tremendous boost for spintronic technologies, EPFL scientists have shown that electrons can jump through spins much faster than previously thought.



Electrons spin around atoms, but also spin around themselves, and can cross over from one spin state to another. A property which can be exploited for next-generation hard drives. However, "spin cross-over" has been considered too slow to be efficient. Using ultrafast measurements, EPFL scientists have now shown for the first time that <u>electrons</u> can cross spins at least 100,000 times faster than previously thought. Aside for its enormous implications for fundamental physics, the finding can also propel the field of spintronics forward. The study is published in *Nature Chemistry*.

The rules of spin

Although difficult to describe in everyday terms, <u>electron spin</u> can be loosely compared to the rotation of a planet or a spinning top around its axis. Electrons can spin in different manners referred to as "spin states" and designated by the numbers 0, 1/2, 1, 3/2, 2 etc. During chemical reactions, electrons can cross from one spin state to another, e.g. from 0 to 1 or 1/2 to 3/2.

Spin cross-over is already used in many technologies, e.g. optical lightemitting devices (OLED), energy conversion systems, and cancer phototherapy. Most prominently, spin cross-over is the basis of the fledgling field of spintronics. The problem is that spin cross-over has been thought to be too slow to be efficient enough in circuits.

Spin cross-over is extremely fast

The lab of Majed Chergui at EPFL has now demonstrated that spin crossover is considerably faster than previously thought. Using the highest <u>time-resolution</u> technology in the world, the lab was able to "see" electrons crossing through four spin states within 50 quadrillionths of a second - or 50 femtoseconds.



"Time resolution has always been a limitation," says Chergui. "Over the years, labs have used techniques that could only measure spin changes to a billionth to a millionth of a second. So they thought that spin cross-over happened in this timeframe."

Chergui's lab focused on materials that show much promise in spintronics applications. In these materials, electrons jump through four <u>spin-states</u>: from 0 to 1 to 2. In 2009, Chergui's lab pushed the boundaries of time resolution to show that this 0-2 "jump" can happen within 150 femtoseconds - suggesting that it was a direct event. Despite this, the community still maintained that such spin cross-overs go through intermediate steps.

But Chergui had his doubts. Working with his postdoc Gerald Auböck, they used the lab's world-recognized expertise in ultrafast spectroscopy to "crank up" the time resolution. Briefly, a laser shines on the material sample under investigation, causing its electrons to move. Another laser measures their spin changes over time in the ultraviolet light range.

The finding essentially demolishes the notion of intermediate steps between spin jumps, as it does not allow enough time for them: only 50 quadrillionths of a second to go from the "0" to the "2" spin state. This is the first study to ever push time resolution to this limit in the ultraviolet domain. "This probably means that it's even faster," says Chergui. "But, more importantly, that it is a direct process."

From observation to explanation

With profound implications for both technology and fundamental physics and chemistry, the study is an observation without an explanation. Chergui believes that the key is electrons shuttling back-andforth between the iron atom at the center of the material's molecules and its surrounding elements. "When the laser light shines on the atom, it



changes the electron's spin angle, affecting the entire spin dynamics in the molecule."

It is now up to theoreticians to develop a new model for ultrafast spin changes. On the experimental side of things, Chergui's lab is now focusing on actually observing electrons shuttling inside the molecules. This will require even more sophisticated approaches, such as core-level spectroscopy. Nonetheless, the study challenges ideas about spin crossover, and might offer long-awaited solutions to the limitations of spintronics.

More information: Auböck G, Chergui M. Sub-50 fs photo-induced spin cross-over in [Fe(bpy)3]2+. *Nature Chemistry* 20 July 2015. DOI: 10.1038/nchem.2305

Provided by Ecole Polytechnique Federale de Lausanne

Citation: Spintronics just got faster (2015, July 20) retrieved 4 May 2024 from <u>https://phys.org/news/2015-07-spintronics-faster.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.