

# Ancient magma movements responsible for Gascoyne minerals

August 20 2015, by Geoff Vivian

---



The deposit's owner, Abra Mining, provided the researchers with drill cores from which they sampled mineralised zones. Credit: Josh

Geologists have used a technique developed at Curtin University to determine magmatic fluids came up from the earth's mantle repeatedly over the past 1600 million years, depositing minerals along a fault line in the Gascoyne region.

Geological Survey of Western Australia researcher Dr Simon Johnson told the Geological Survey Open Day earlier this year the fault had been active 1600, 1375, 1220 and 1000 million years ago.

"Each time that fault gets reactivated there are hydrothermal fluids which percolate through that rock system—they cause alterations and they cause alteration of the ore deposit," he said.

He said the usual analysis method, by which a rock sample was crushed before analysis, had made it impossible to date when the various minerals present had first been deposited.

However, Professor Birger Rasumussen's technique, which involves mounting extremely thin slices of rock on a slide, meant small mineral crystals could be identified and individually dated in situ.

Dr Johnson says they applied the technique to a diamond drill core from the Abra deposit.

This is the biggest known [mineral deposit](#) in the Capricorn Orogen, which is a deep margin between two former [tectonic plates](#).

He said Abra is a huge lead-zinc deposit, with accessory gold and silver, which is completely covered by younger rocks.

It was detected in a magnetic survey, confirmed by drilling in the early 1980s, and is now a mining lease.

## **Samples dished up as ultra thin disks**

The deposit's owner, Abra Mining, provided the researchers with drill cores from which they sampled mineralised zones.

Using Prof Rasmussen's technique they, inspecting them for phosphate minerals like xenotime or monazite which they drilled out as 3mm disks.

They mounted these on tape, encased it in an epoxy glue and placed it in a Sensitive High Resolution Ion Micro Probe (SHRIMP) machine.

This allowed them to fire [oxygen ions](#) at the grains of xenotime or the monazite.

The oxygen ions evaporate part of the xenotime or the monazite which they fed into a Faraday collector.

This dated the xenotime grains by measuring various isotope ratios of uranium, thorium, lead and intermediate products.

While it sits at the junction of the former Yilgarn and Pilbara tectonic plates, both of which are rich in minerals, the orogeny has very few working mines.

Dr Johnson said two Curtin University PhD students will apply the new method in an attempt to locate more prospective areas of the orogeny for mineral explorers.

Provided by Science Network WA

Citation: Ancient magma movements responsible for Gascoyne minerals (2015, August 20) retrieved 12 May 2024 from <https://phys.org/news/2015-08-ancient-magma-movements->

[responsible-gascoyne.html](#)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.