


The consortium of motile and cellulose degrading bacteria can be used for solid state cellulose hydrolysis

August 4 2015



Two colonies of *Paenibacillus vortex* are shown. White round structures at the edges of the colony called the vortices and are responsible for cargo carriage. The colors were inverted for better contrast. Scale bar -9mm. (b) Schematic presentation of *P. vortex* and *E. coli* collaboration. *P. vortex* can move on solid surfaces, carry *E. coli* while *E. coli* contributes its cellulases. During the collaboration, the underlying colony of *P. vortex* is constantly moving on the solid surface (plant biomass) that contains cellulose. The consortium allows

microorganism distribution and multi-enzymatic biomass hydrolysis. The *E. coli* cells (with pCellulose plasmid) are transported by *P. vortex*. During the transport, *E. coli* degrades cellulose into monomers and dimers, making it available for uptake by both microorganisms. Credit: *TECHNOLOGY*

Cellulose hydrolysis has many industrial applications in such fields as biofuel production, food, paper, cosmetic, pharmaceutical industries and textile manufacture. A novel approach to cellulose hydrolysis using a consortium of motile bacteria moving on solid surfaces and carrying microbial lug-gage - another bacteria that can efficiently hydrolyze cellulose, was demonstrated by the group of Professor Eshel Ben- Jacob (R.I.P) from Tel Aviv University.

Cellulose is the most abundant polysaccharide on earth, however, it is also one of the hardest to hydrolyze. Currently, the conversion of cellulosic materials to renewable energy includes either chemical or biological [hydrolysis](#) followed by fermentation of sugars to ethanol/butanol. Utility cost of biological- [enzymatic hydrolysis](#) is low compared to chemical hydrolysis since it is usually conducted at mild conditions and does not have a corrosion problem. Biological hydrolysis is usually carried out in liquid. However, Solid State hydrolysis is financially feasible due to lower capital investment and lower operating expenses. Most cellulolytic bacteria are anaerobes with a very low growth rate and low enzyme titers.

Current research utilized a recombinant strain of *E. coli*, aerobic bacteria with high growth rate and enzyme titers. The *E. coli* contains α -glucosidase and cellulase genes (on pCellulose plasmid constructed in the lab of Prof. Jay Keasling). The un-motile *E. coli* was shown to efficiently degrade cellulose both in liquid and solid mediums.

The innovation of the approach is the use of the recombinant *E. coli* as a microbial luggage of another highly [motile bacteria](#)- *Paenibacillus vortex*. *Paenibacillus vortex* is known for its ability to swarm (coordinated migration on solid surfaces using flagella) and collaborate with other microorganisms by carrying them in specialized structures called vortices (Fig 1 a). Thus, two bacterial species create a mutually beneficial consortium in which *Paenibacillus vortex* provides a motile force while the *E. coli* degrades cellulose into monomers and dimers, making it available for uptake by both microorganisms.

Moreover, *Paenibacillus vortex* has the ability to degrade xylan, thus contributing its own enzymes to the consortium making it even more effective for degradation of the plant material. Since *Paenibacillus vortex* has the ability to swarm on solid mediums it might be relevant to use the consortium in Solid State hydrolysis of plant materials such as crops leftovers. This application of using crops needs more research, and, maybe, addition of other microorganisms relevant for fermentation. If it is be feasible it would allow easier and cheaper way to hydrolyze [cellulose](#).

More information: Mark Polikovsky, Eshel Ben-Jacob, and Alin Finkelshtein, *Technology* [DOI: 10.1142/S2339547815400087](https://doi.org/10.1142/S2339547815400087)

Provided by World Scientific Publishing

Citation: The consortium of motile and cellulose degrading bacteria can be used for solid state cellulose hydrolysis (2015, August 4) retrieved 18 April 2024 from <https://phys.org/news/2015-08-consortium-motile-cellulose-degrading-bacteria.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.