


Another milestone in hybrid artificial photosynthesis

August 25 2015, by Lynn Yarris



A team of researchers at the U.S. Department of Energy (DOE)'s Lawrence Berkeley National Laboratory developing a bioinorganic hybrid approach to artificial photosynthesis have achieved another milestone. Having generated quite a buzz with their hybrid system of semiconducting nanowires and bacteria that used electrons to synthesize carbon dioxide into acetate, the team has now developed a hybrid system that produces renewable molecular hydrogen and uses it to synthesize carbon dioxide into methane, the primary constituent of natural gas.

"This study represents another key breakthrough in solar-to-chemical [energy conversion efficiency](#) and [artificial photosynthesis](#)," says Peidong Yang, a chemist with Berkeley Lab's Materials Sciences Division and one of the leaders of this study. "By generating [renewable hydrogen](#) and feeding it to microbes for the production of methane, we can now expect an electrical-to-chemical efficiency of better than 50 percent and a solar-to-chemical [energy](#) conversion efficiency of 10-percent if our system is coupled with state-of-art solar panel and electrolyzer."

Yang, who also holds appointments with UC Berkeley and the Kavli Energy NanoScience Institute (Kavli-ENSI) at Berkeley, is one of three corresponding authors of a paper describing this research in the Proceedings of the National Academy of Sciences (PNAS). The paper is titled "A hybrid bioinorganic approach to solar-to-chemical conversion." The other corresponding authors are Michelle Chang and Christopher Chang. Both also hold joint appointments with Berkeley Lab and UC Berkeley. In addition, Chris Chang is a Howard Hughes Medical Institute (HHMI) investigator.

Photosynthesis is the process by which nature harvests the energy in sunlight and uses it to synthesize carbohydrates from [carbon dioxide](#) and water. Carbohydrates are biomolecules that store the chemical energy used by living cells. In the original hybrid artificial photosynthesis system developed by the Berkeley Lab team, an array of silicon and titanium oxide nanowires collected solar energy and delivered electrons to microbes which used them to reduce carbon dioxide into a variety of value-added chemical products. In the new system, solar energy is used to split the water molecule into molecular oxygen and [hydrogen](#). The hydrogen is then transported to microbes that use it to reduce carbon dioxide into one specific chemical product, methane.



A major advance in artificial photosynthesis poses win/win for the environment – removing CO₂ from the atmosphere and using it for green chemistry, including renewable fuel production. Credit: Caitlin Givens

"In our latest work, we've demonstrated two key advances," says Chris Chang. "First, our use of renewable hydrogen for [carbon dioxide fixation](#) opens up the possibility of using hydrogen that comes from any sustainable energy source, including wind, hydrothermal and nuclear. Second, having demonstrated one promising organism for using renewable hydrogen, we can now, through synthetic biology, expand to other organisms and other value-added chemical products."

The concept in the two studies is essentially the same – a membrane of semiconductor nanowires that can harness solar energy is populated with bacterium that can feed off this energy and use it to produce a targeted

carbon-based chemical. In the new study, the membrane consisted of indium phosphide photocathodes and titanium dioxide photoanodes. Whereas in the first study, the team worked with *Sporomusa ovata*, an anaerobic bacterium that readily accepts electrons from the surrounding environment to reduce carbon dioxide, in the new study the team populated the membrane with *Methanosarcina barkeri*, an anaerobic archaeon that reduces carbon dioxide using hydrogen rather than electrons.

"Using hydrogen as the energy carrier rather than electrons makes for a much more efficient process as molecular hydrogen, through its chemical bonds, has a much higher density for storing and transporting energy," says Michelle Chang.

In the newest membrane reported by the Berkeley team, [solar energy](#) is absorbed and used to generate hydrogen from water via the hydrogen evolution reaction (HER). The HER is catalyzed by earth-abundant nickel sulfide nanoparticles that operate effectively under biologically compatible conditions. Hydrogen produced in the HER is directly utilized by the *Methanosarcina barkeri* archaeons in the membrane to produce methane.

"We selected methane as an initial target owing to the ease of product separation, the potential for integration into existing infrastructures for the delivery and use of natural gas, and the fact that direct conversion of carbon dioxide to methane with synthetic catalysts has proven to be a formidable challenge," says Chris Chang. "Since we still get the majority of our methane from [natural gas](#), a fossil fuel, often from fracking, the ability to generate methane from a renewable hydrogen source is another important advance."

Adds Yang, "While we were inspired by the process of natural photosynthesis and continue to learn from it, by adding nanotechnology

to help improve the efficiency of natural systems we are showing that sometimes we can do even better than nature."

More information: "Hybrid bioinorganic approach to solar-to-chemical conversion." *PNAS* 2015 ; published ahead of print August 24, 2015, [DOI: 10.1073/pnas.1508075112](https://doi.org/10.1073/pnas.1508075112)

Provided by Lawrence Berkeley National Laboratory

Citation: Another milestone in hybrid artificial photosynthesis (2015, August 25) retrieved 29 April 2024 from <https://phys.org/news/2015-08-milestone-hybrid-artificial-photosynthesis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.