


Peat fires in Sumatra strengthen in El Nino years

August 21 2015, by Lynn Jenner



NASA's Aqua satellite collected this natural-color image with the Moderate

Resolution Imaging Spectroradiometer, MODIS, instrument on June 05, 2013. Actively burning areas, detected by MODIS's thermal bands, are outlined in red. NASA image courtesy Jeff Schmaltz LANCE/EOSDIS MODIS Rapid Response Team, GSFC. Caption by Lynn Jenner Credit: Jeff Schmaltz LANCE/EOSDIS MODIS Rapid Response Team, GSFC.

Sumatra's burning again and the El Nino event that is occurring this year is partially to blame for the proliferation of the blazes.

El Nino events bring drier weather to Indonesia and with it, a rise in the number of [forest fires](#) that break out. Many of the wildfires occur on land that has deep reserves of peat, which is rich in carbon. These fires produce huge plumes of dark smoke, insidious carbon dioxide, and other [greenhouse emissions](#) that contribute to the current climate change crisis.

Sumatra, Indonesia might seem a world away but the smoke from the fires there tends to travel along the jetstreams bringing that smoke and its accompanying degraded air quality across the globe.

Many of the fires on the island are due to farmers clearing their land for agricultural reasons. These fires which can usually be contained outgrow their boundaries and become raging infernos more often during the drier seasons. These types of fires are illegal in Sumatra but are still used by the subsistence farmers as they have for thousands of years.

Provided by NASA's Goddard Space Flight Center

Citation: Peat fires in Sumatra strengthen in El Nino years (2015, August 21) retrieved 20 March 2024 from <https://phys.org/news/2015-08-peat-sumatra-el-nino-years.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.