


Snaring a dark energy 'chameleon': Is dark energy hiding from us? Matter may be screening it from our view

August 20 2015



UC Berkeley physicists used an atom interferometer to try to detect anomalous forces from a chameleon field, a hypothetical dark energy particle. If chameleons exist, they would have a very small effect on the interaction between cesium atoms and an aluminum sphere. The group's failure to detect this effect narrows the search for the particle. Credit: Holger Muller/UC Berkeley

If dark energy is hiding in our midst in the form of hypothetical particles called "chameleons," Holger Müller and his team at the University of California, Berkeley, plan to flush them out.

The results of an experiment reported in this week's issue of *Science* narrows the search for chameleons a thousand times compared to previous tests, and Müller, an assistant professor of physics, hopes that his next experiment will either expose chameleons or similar ultralight particles as the real dark energy, or prove they were a will-o'-the-wisp after all.


Dark energy was first discovered in 1998 when scientists observed that the universe was expanding at an ever increasing rate, apparently pushed apart by an unseen pressure permeating all of space and making up about 68 percent of the energy in the cosmos. Several UC Berkeley scientists were members of the two teams that made that Nobel Prize-winning discovery, and physicist Saul Perlmutter shared the prize.

Since then, theorists have proposed numerous theories to explain the still mysterious energy. It could be simply woven into the fabric of the universe, a cosmological constant that Albert Einstein proposed in the equations of general relativity and then disavowed. Or it could be quintessence, represented by any number of [hypothetical particles](#), including offspring of the Higgs boson.

In 2004, theorist and co-author Justin Khoury of the University of Pennsylvania proposed one possible reason why dark energy particles haven't been detected: they're hiding from us.

Specifically, Khoury proposed that dark energy particles, which he dubbed chameleons, vary in mass depending on the density of surrounding matter.

In the emptiness of space, chameleons would have a small mass and exert force over long distances, able to push space apart. In a laboratory, however, with matter all around, they would have a large mass and extremely small reach. In physics, a low mass implies a long-range force, while a high mass implies a short-range force.



The vacuum chamber of the atom interferometer contains a one-inch diameter aluminum sphere. If chameleons exist, cesium atoms would fall toward the sphere with a slightly greater acceleration than their gravitational attraction would predict. Credit: Holger Muller

This would be one way to explain why the energy that dominates the universe is hard to detect in a lab.

"The chameleon field is light in empty space but as soon as it enters an object it becomes very heavy and so couples only to the outermost layer of a big object, and not to the internal parts," said Müller, who is also a faculty scientist at Lawrence Berkeley National Laboratory. "It would pull only on the outermost nanometer."

Lifting the camouflage

When UC Berkeley post-doctoral fellow Paul Hamilton read an article by theorist Clare Burrage last August outlining a way to detect such a particle, he suspected that the atom interferometer he and Müller had built at UC Berkeley would be able to detect chameleons if they existed. Müller and his team have built some of the most sensitive detectors of forces anywhere, using them to search for slight gravitational anomalies that would indicate a problem with Einstein's General Theory of Relativity. While the most sensitive of these are physically too large to sense the short-range chameleon force, the team immediately realized that one of their less sensitive atom interferometers would be ideal.

Burrage suggested measuring the attraction caused by the chameleon field between an atom and a larger mass, instead of the attraction between two large masses, which would suppress the chameleon field to the point of being undetectable.

That's what Hamilton, Müller and his team did. They dropped cesium atoms above an inch-diameter aluminum sphere and used sensitive lasers to measure the forces on the atoms as they were in free fall for about 10 to 20 milliseconds. They detected no force other than Earth's gravity, which rules out chameleon-induced forces a million times weaker than gravity. This eliminates a large range of possible energies for the

particle.

What about symmetrons?

Experiments at CERN in Geneva and the Fermi National Accelerator Laboratory in Illinois, as well as other tests using neutron interferometers, also are searching for evidence of chameleons, so far without luck. Müller and his team are currently improving their experiment to rule out all other possible particle energies or, in the best-case scenario, discover evidence that chameleons really do exist.

"Holger has ruled out chameleons that interact with normal matter more strongly than gravity, but he is now pushing his experiment into areas where chameleons interact on the same scale as gravity, where they are more likely to exist," Khoury said.

Their experiments may also help narrow the search for other hypothetical screened dark energy fields, such as symmetrons and forms of modified gravity, such as so-called $f(R)$ gravity.

"In the worst case, we will learn more of what [dark energy](#) is not. Hopefully, that gives us a better idea of what it might be," Müller said. "One day, someone will be lucky and find it."

More information: "Atom-interferometry constraints on dark energy," by P. Hamilton et al: www.sciencemag.org/lookup/doi/10.1126/science.aaa8883

Provided by University of California - Berkeley

Citation: Snaring a dark energy 'chameleon': Is dark energy hiding from us? Matter may be screening it from our view (2015, August 20) retrieved 9 September 2024 from <https://phys.org/news/2015-08-snaring-dark-energy-chameleon-screening.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.