

# Widest distribution of mammoths during the last Ice Age

August 25 2015

---



Prof. R.-D. Kahlke inspects the newly preserved femur of a mammoth. Credit: T. Korn/Senckenberg Weimar

Ice Age paleontologist Prof. Dr. Ralf-Dietrich Kahlke of the Senckenberg Research Station for Quaternary Paleontology in Weimar

recorded the maximum geographic distribution of the woolly mammoth during the last Ice Age and published the most accurate global map in this regard. The ice-age pachyderms populated a total area of 33,301,000 square kilometers and may thus be called the most successful large mammals of this era. The study, recently published online in the scientific journal *Quaternary International*, determined that the distribution was limited by a number of climate-driven as well as climate-independent factors.

The mammoth is the quintessential symbol of the Ice Age – and the status of these shaggy pachyderms has now been confirmed scientifically. "The recent research findings show that during the last Ice Age, mammoths were the most widely distributed large mammals, thus rightfully serving as a flagship species of the glacial era," according to Prof. Dr. Ralf-Dietrich Kahlke, an Ice Age researcher at the Senckenberg Research Station for Quaternary Paleontology in Weimar.

Kahlke has summarized the [mammoth](#)'s distribution during the most recent Ice Age, i.e., the period between approx. 110,000 and 12,000 years ago, on a worldwide map. All in all, the Weimar paleontologist determined a total distribution area of 33,301,000 square kilometers for these large mammals – almost 100 times the area of Germany today. From Portugal in the southwest across Central and Eastern Europe, Mongolia, Northern China, South Korea and Japan up to Northeastern Siberia, and thence to the American Midwest and Eastern Canada, from the shelf regions of the Arctic Ocean and Northwestern Europe to the bottom of the Adriatic Sea and to the mountains of Crimea: the fossil remains of woolly mammoths have been found everywhere.

"We related the computed distribution area to the real land surface at that time, thus generating the most precise map to date regarding the global habitats of the [woolly mammoth](#)," explains Kahlke, and he adds, "Such detailed knowledge regarding the distribution area is not even

available for many species of animals alive today."

The generated map is based on decades of surveys of thousands of excavation sites on three continents. "Even sites under water, off the North American Atlantic shore and the North Sea, were taken into account. Due to the lower sea levels during the Ice Age – a large volume of water was bound in glaciers – these areas had fallen dry and were also inhabited by *Mammuthus primigenius*," according to Kahlke.





Complete left tusk of an ice-age Woolly Mammoth (*Mammuthus primigenius*) from the Siberian Arctic on the Taimyr Peninsula. Each individual discovery increases our knowledge about the past distribution of these Ice Age giants. Credit: R.-D. Kahlke/ Senckenberg Weimar The Siberian permafrost soil contains the best preserved mammoth remains in the world. A recently discovered tusk is carried off for inspection. Credit: R.-D. Kahlke/ Senckenberg Weimar

Only the [ice-age](#) bison (*Bison priscus*) had a widespread distribution similar to that of the mammoths. Kahlke explains, "The bison were clearly more variable than the woolly mammoths. Obviously, the mammoths had a higher tolerance toward various environmental factors and they were able to successfully settle in a variety of rather different open landscapes."

But there were certain factors that limited the distribution of the hirsute pachyderms: glaciers, mountain chains, semi-deserts and deserts, as well as changes in sea level and shifts in vegetation placed restrictions on the mammoths' distribution area. "The analysis of these limiting factors is useful in understanding the distribution of fossil species and their extinction – as with the mammoths toward the end of the last Ice Age. In addition, the data aid in comprehending current changes in the distribution areas of recent animal species," offers Kahlke in summary.



Complete left tusk of an ice-age Woolly Mammoth (*Mammuthus primigenius*) from the Siberian Arctic on the Taimyr Peninsula. Each individual discovery increases our knowledge about the past distribution of these Ice Age giants.  
Credit: R.-D. Kahlke/ Senckenberg Weimar

**More information:** "The maximum geographic extension of Late Pleistocene *Mammuthus primigenius* (Proboscidea, Mammalia) and its limiting factors," *Quaternary International* (2015), [dx.doi.org/10.1016/j.quaint.2015.03.023](https://doi.org/10.1016/j.quaint.2015.03.023)

Provided by Senckenberg Research Institute and Natural History  
Museum

Citation: Widest distribution of mammoths during the last Ice Age (2015, August 25) retrieved 5  
May 2024 from <https://phys.org/news/2015-08-widest-mammoths-ice-age.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.