

Plant debris decomposition tied to manganese

September 29 2015



Lab researchers have shown that manganese plays a strong role in plant debris decomposition in forest ecosystems.

The decomposition of plant debris (litter) is a fundamental process that regulates the release of nutrients for plant growth and the formation of soil organic matter in forest ecosystems.

A strong correlation has previously been observed between litter manganese (Mn) content and decomposition rates across a variety of [forest ecosystems](#). However, the mechanisms underlying Mn's role in litter decomposition were not well understood. Until now.

In a recent article in *Proceedings of the National Academy of Sciences* (PNAS), Livermore Graduate Scholar Marco Keiluweit and LLNL scientist Jennifer Pett-Ridge show that long-term litter decomposition rate in forest ecosystems is tightly coupled to manganese (Mn) redox cycling. (Redox reactions include all chemical reactions in which atoms have their oxidation state changed).

The work has implications to the understanding and prediction of carbon cycling in [terrestrial ecosystems](#).

Litter decomposition rates are strongly influenced by climatic factors such as temperature and moisture, but also are linked to litter chemistry.

The team's results suggest that the litter-decomposing machinery in the coniferous forest site they studied depends on the ability of plants and microbes to supply, accumulate and regenerate short-lived Mn species in the litter layer.

"This implies that the bioavailability, mobility and reactivity of Mn in the plant–soil system have a profound impact on litter decomposition rates, and therefore the overall global carbon cycle," Pett-Ridge said.

Over seven years of litter decomposition, the observed microbial transformation was strongly correlated to variations in Mn oxidation states and concentration. A detailed chemical imaging analysis of the litter revealed that soil fungi incorporate unreactive Mn provided by fresh plant litter to produce and redistribute reactive (oxidizing) Mn species at sites of active litter decay. Formation of reactive Mn species

coincided with the generation of oxidation breakdown products of aromatic organic compounds, providing direct proof of the role of Mn-based oxidizers in the breakdown of litter.

"Incorporating the coupling of litter decomposition and other elemental cycles into conceptual and numerical models may significantly improve our mechanistic understanding and prediction of carbon cycling in terrestrial ecosystems," Pett-Ridge said.

More information: "Long-term litter decomposition controlled by manganese redox cycling." *PNAS* 2015 112 (38) E5253-E5260; published ahead of print September 8, 2015, [DOI: 10.1073/pnas.1508945112](https://doi.org/10.1073/pnas.1508945112)

Provided by Lawrence Livermore National Laboratory

Citation: Plant debris decomposition tied to manganese (2015, September 29) retrieved 2 May 2024 from <https://phys.org/news/2015-09-debris-decomposition-tied-manganese.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.