

# Evidence for solid-liquid critical points of water in carbon nanotubes

September 24 2015




Figure1: Hydrogen bond structures of six ices formed in the nanotubes: (a) (4,0)ice, (b) (5,0) ice and (c) filled (6,0) ice at a diameter of 1.1nm, (d) (6,0) ice, (e) filled (7,0) ice and (f) filled (8,0) ice at a diameter of 1.25nm. Top views and the corresponding side view are drawn abreast. Central water molecules forming a chain in the filled ices are colored red to distinguish them from the

exterior rings.

Many physicists do not accept the idea that a solid-liquid phase boundary can terminate at a critical point—a unique state where two phases lose their separate identities. Why do they not? The authoritative textbook by Landau and Lifshitz says that "we can say only that a particular symmetry property exists or does not exist; ... The critical point therefore cannot exist for such phases." But since 2001, the possibility of the solid-liquid critical point has been reported in computer-simulation studies of water in nanopores. In fact, there is no rigorous proof for the nonexistence of the solid-liquid critical point.

Kenji Mochizuki and Kenichiro Koga at Okayama University provided unambiguous evidence to support the solid-liquid critical point for a class of [water](#) in nanotubes by performing extensive molecular dynamics simulations: macroscopic solid-liquid phase separation below a critical temperature  $T_c$ , diverging heat capacity and isothermal compressibility at around  $T_c$ , and the loci of response function maxima (the Widom lines) above  $T_c$ .

Figure 1 shows the hydrogen bond structures of six crystals of ice formed in carbon nanotubes with diameters of 1.11 nm and 1.25 nm. The researchers found that all of the first-order phase boundaries between the ice and liquid eventually ceased to exist at the critical points, as shown in Figure 2. The T-P phase diagram is very different from that of bulk water, where ice regions are completely surrounded by the first-order phase boundaries (solid black lines).

The researchers also gave a microscopic explanation to a simple, yet unanswered, question: how can liquid water continuously freeze to crystalline ice? They found dynamic fluctuations of microscopic

domains of water and [ice](#) near the critical point—a microscopic picture of water in the course of gradual freezing or melting.

The critical points in confined water are ubiquitous and can be found at ambient conditions by tuning the pore diameter, and therefore one of the potential applications is to use the tunable critical fluctuations in order to facilitate chemical reactions, structural changes in biological molecules, and formation of biomolecules assembly in water.




Figure 2: Phase diagrams of (a) bulk water, (b) water confined in the nanotube of a diameter of 1.11 nm, and (c) 1.25 nm. The phase diagrams of (b) and (c) are firstly revealed in this study. Solid regions are colored in blue and critical points are indicated by red circles.

**More information:** "Solid-liquid critical behavior of water in nanopores." *Proc. Natl. Acad. Sci. USA* (2015) (DOI): 10.1073/pnas.1422829112

Provided by Okayama University

Citation: Evidence for solid-liquid critical points of water in carbon nanotubes (2015, September 24) retrieved 2 May 2024 from <https://phys.org/news/2015-09-evidence-solid-liquid-critical-carbon-nanotubes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.